

**Research Article**

## The potential use of humic acid-coated biochar for reducing Pb and Cu in the soil to improve plant growth

Amir Hamzah<sup>1</sup>, Rosyda Priyadarshini<sup>2\*</sup>, Astuti<sup>1</sup>

<sup>1</sup> Tribhuwana Tunggal University, Jl Telaga Warna, Tlogomas, Malang 65144, East Java, Indonesia

<sup>2</sup> University of Pembangunan Nasional Veteran Jawa Timur, Jl Raya Rungkut Madya Gunung Anyar, Surabaya 60294, East Java, Indonesia

\*corresponding author: rosyda\_p@upnjatim.ac.id

### Abstract

#### Article history:

Received 22 May 2022

Accepted 13 September 2022

Published 1 October 2022

#### Keywords:

humic acid-coated biochar

heavy metals

plant growth

The use of biochar for remediating heavy metal-polluted soils is still partial. Various methods of controlling soil pollution are currently being implemented by combining several methods. One of which is coating the biochar with humic acid to increase the effectiveness of nutrient uptake by plants. This study aimed to elucidate the effect of humic acid-coated biochar (HCB) on reducing Pb and Cu in the soil to improve plant growth. Treatments tested were combinations of two factors. The first factor was the dose of HCB, namely D0 = control (without HCB), D1 = 15 t HCB ha<sup>-1</sup>, D2 = 30 t HCB ha<sup>-1</sup>, and D3 = 45 t HCB ha<sup>-1</sup>. The second factor was the type of plants, namely spinach (P1), water spinach (P2) and mustard green (P3). Twelve treatments were arranged in a randomized block design with three replications. The parameters observed were plant height, plant leaf area, plant stem diameter, and plant fresh weight. The results showed that the best plant growth was achieved at a dose of 30 t HCB ha<sup>-1</sup>. The mustard green had the highest Pb uptake (0.025 g pot<sup>-1</sup>), and the lowest Pb uptake (0.014 g pot<sup>-1</sup>) was observed for water spinach. The highest Cu uptake (0.443 g pot<sup>-1</sup>) was observed in water spinach, followed by spinach (0.282 g pot<sup>-1</sup>) and mustard green (0.143 g pot<sup>-1</sup>). In general, the amount of Pb reduced by plants ranged from 40.04 to 87.28%, and the amount of Cu by plants ranged from 8.63 to 40.23%.

**To cite this article:** Hamzah, A., Priyadarshini, R. and Astuti. 2022. The potential use of humic acid-coated biochar for reducing Pb and Cu in the soil to improve plant growth. *Journal of Degraded and Mining Lands Management* 10(1):4001-4009, doi:10.15243/jdmlm.2022.101.4001.

### Introduction

Heavy metal contamination will affect agricultural production, human health, environmental problems, and food quality throughout the food chain. Several countries, such as Japan, China, Korea, Thailand, and Indonesia, have been contaminated by heavy metals (Hamzah et al., 2017). Some metals are essential, but certain metals, such as Cd and Pb, are toxic and dangerous for human health. Some countries have exceeded the threshold value of heavy metals level. Soil Pb level in China is more than 0.2 mg kg<sup>-1</sup> (Shifaw, 2018), while the Cd level in horticultural centres in East Java has reached 2.26 mg kg<sup>-1</sup> (Hamzah

et al., 2016; 2017). Heavy metal accumulation under agricultural land will affect the soil physical, biological, and chemical properties (Satpathy et al., 2014); the heavy metal will enter the plant root, leaves, and grain (Singh et al., 2010). Wang (2005).

Plants have different ways of absorbing and accumulating heavy metals in their edible parts; that can directly transfer the heavy metals to the human body. The soil-to-plant transfer of heavy metals is of great importance (Kalavrouziotis et al., 2011). Heavy metals uptake by plant roots and their transfer and accumulation in soil rice have become interesting to study. Among agricultural products, rice is widely

consumed as staple food worldwide, especially in Asia. In Asia, rice was the most source of soil Cd and Pb contaminants for human health (Zhang and Ke, 2004), especially in Japan (Tsukahara et al., 2003). Other agricultural products like horticulture products are also threatened with heavy metal contamination. Therefore, how to manage, stabilize, and immobilize the heavy metal is being concerned.

Biochar is carbon-rich material produced through the pyrolysis process at high temperatures with limited oxygen supply (Ahmad et al., 2013; Lu et al., 2014). Biochar was proved very effective in improving soil properties and crop yield (Jeffery et al., 2011). Biochar has good adsorption capacity and the potential as a carrier for slow-release fertilizer (Qin et al., 2016). Biochar is also known to be a reasonably effective solution to improve soil physical properties in the short term, but new chemical properties will be seen in a long time (Lehman et al., 2011). Many studies showed the capability of biochar in immobilizing trace elements (e.g. Cu), minimizing the phytotoxicity and detrimental effects of Cu (Park et al., 2016). According to Lu et al. (2014), the addition of biochar could increase the above-ground biomass and soil pH but lower the solubility of heavy metals. This is in line with the reduction of Cu accumulated in plant biomass. Biochar has good adsorption capacity and the potential as a carrier for slow-release fertilizer (Qin et al., 2016). The utilization of biochar is very effective for improving soil physical characteristics in a short period, and in the long term, the soil chemical characteristics will also improve (Lehman et al., 2011).

Humic substances are well known as one of the amendments used for the phytoextraction of heavy metals from soil. The humic substance has a complex molecular structure and many functional groups that are useful in stabilizing and immobilizing metal ions, especially cation. Humic acid will reduce the metal bioavailability and increase its capability in chelating metal ions with their functional groups. Humic acid affects heavy metal availability and mobility of heavy metals (Rong et al., 2020). Humic acid and biochar will transform the exchangeable fraction into the non-exchangeable fraction (Zhou et al., 2018) due to their large area and the high content of organic functional groups (Liu et al., 2022).

The purpose of the remediation method is to suppress the availability of heavy metals so that they would be difficult to absorb by plants. Humic acid stimulates and activates biological and physiological processes for microbes that live and nest in the soil and act as ameliorants (Ouni et al., 2014). Humic acid will also stimulate soil microbial activity in providing soil nutrients. Liu et al. (2022) observed that functional groups of humic substances require a suitable pH for binding heavy metals. The humic substance shows a good effect on the complexation. It decreases heavy metal bioavailability, but a single application of humic substances has a poor impact on available heavy metals in paddy plant tissues. Therefore, it is necessary

to develop a new remediation scheme for heavy metal remediation by coupling humic substance and biochar.

## Materials and Methods

### *Preparation of humic acid-coated biochar*

Humic acid-coated biochar (HCB) is a mixture of biochar, humic acid, and adhesive polymer. The biochar was made from the rice straw by pyrolysis combustion at a temperature of 400 °C, while the humic acid was extracted from the organic matter using a modified method by Gayathri et al. (2019). Ten grams of organic matter was extracted with 100 ml of 0.5 N NaOH (1:10), shaken for 24 hours, cooled for 16 hours, centrifuged at 1500 rpm, and then filtered again to obtain humic compounds. These substances were added with 6 N HCl until the pH of the solution reached 2 and separated again with filter paper. The precipitate obtained was rinsed with distilled water to remove residual chloride. Then, it was oven-dried at 105 °C to characterize and determine the percentage of humic acid. The HCB was made by thoroughly mixing biochar and humic acid in a balanced ratio. One gram of adhesive polymer was added to the mixture.

### *Pot experiment*

A pot experiment was conducted from April to July 2021, in Malang, East Java, at an altitude of  $\pm 450$  m above sea level. Treatments tested were combinations of two factors. The first factor was the dose of HCB, i.e. D0 = Control (without HCB), D1 = 15 t HCB ha<sup>-1</sup>, D2 = 30 t HCB ha<sup>-1</sup>, and D3 = 45 t HCB ha<sup>-1</sup>. The second factor was the type of plants, i.e. spinach (P1), water spinach (P2) and mustard green (P3). Twelve treatments were arranged in a randomized block design with three replications. The HCB was applied to 5 kg of soil and incubated for one week before planting. The soil used in this experiment was air-dried topsoil from the rice field at Karangploso village, Malang Regency, East Java. The soil has the following characteristics: pH = 6.34 (neutral), organic C = 2.66% (medium), total N = 0.27% (medium), K<sub>2</sub>O = 0.725 cmol kg<sup>-1</sup> (very low), P<sub>2</sub>O<sub>5</sub> = 58.77 mg kg<sup>-1</sup> (high), Pb = 13.73 mg kg<sup>-1</sup> (critical value), and Cu = 80.23 mg kg<sup>-1</sup> (critical value). Soil and plant analysis were carried out in the laboratory of the Faculty of Agriculture, Tribhuwana Tungadewi University, and the laboratory of the Soil Department, University of Pembangunan Nasional Veteran, East Java.

Plant height, plant leaf area, and plant stem diameter were measured every week after transplanting (WAT) until the harvesting time of 8 WAT. The plant growth measurement was made on three samples per treatment. The leaf area was measured using the constant method (Susilo, 2016). At the end of the experiment (8 WAT), the plant parts were separated into roots, stems, and leaves. The plant parts were washed with running water to remove impurities, and they were then dried in an oven at 60 °C for 72 hours for Pb and Cu analyses.

### Pb and Cu uptake by plants

Pb and Cu uptake by plants was calculated by multiplying the percentage of Pb and Cu concentrations in plant parts with the dry weight of the plant parts. Pb and Cu contents in the plant parts were determined by weighing 2 g of the ground plant sample and then dissolving the sample with 10 mL of HNO<sub>3</sub> and HClO<sub>4</sub> and heating until the remaining volume was 2 mL. It was gradually reheated with distilled water until the liquid became clear (clean white). The clear liquid was then mixed with distilled water and filtered by the 3050 EPA method. The results of the filtering were then measured for the Cu and Pb content using an atomic absorption spectrophotometer based on light absorption with a wavelength range of 190-800 nm (Saputo et al., 2016; Maurya et al., 2018).

### Statistical analysis

The data obtained were subjected to the one-way analysis of variance (ANOVA). The Least Significant

Difference (LSD) test at a significant level of  $p < 0.05$  was performed to see the effect of humic acid-coated biochar doses and plant types on Pb and Cu uptake by the plants and plant growth.

## Results and Discussion

### Plant growth

Spinach, water spinach, and mustard green plants responded positively to the application of humic acid-coated biochar (HCB) in the soil containing critical levels of Pb and Cu. This can be seen from the increase in plant height presented in Figure 1. The application of 30 t HCB ha<sup>-1</sup> gave the tallest plant throughout, while the pot experiment with no HCB applied gave the shortest plant. The average plant weight was 36-45 cm, 35-46 cm, and 21-28 cm for water spinach, spinach, and mustard green, respectively. This indicates that the HCB treatment is effective for improving plant growth.




Figure 1. Effect of HCB application on plant height: (a) water spinach, (b) spinach, and (c) mustard green. D0 = control, D1 = 15 t HCB ha<sup>-1</sup>, D2 = 30 t HCB ha<sup>-1</sup>, D3 = 45 t HCB ha<sup>-1</sup>.

Plant and shoot growth are affected by functional groups of humic acid, i.e. carboxylic (COOH) and phenolic (OH) groups (De Melo et al., 2016). Humic acid is important in enhancing nutrient uptake; thus, plant height increases. Improvement of nutrient uptake (especially N, P, and S) by the humic acid activity has been reported by Atiyeh et al. (2002) and Arancon et

al. (2003). Acceleration of nutrient uptake enhances nitrogen metabolism and protein production, which ultimately increases the chlorophyll contents (Haghighi et al., 2012). This is also related to the humic acid function in increasing membrane cell permeability, photosynthesis, phosphate uptake, and root elongation (Ertani et al., 2015). The application of

30 t HCB ha<sup>-1</sup> also significantly increased the leaf area of water spinach, mustard green, and spinach plants (Figure 2). The leaf area of the plant with no HCB added (D0) area was the smallest for the plant height (Figure 1). Humic compounds can increase vegetative growth presumably due to their capability to increase both the soil chemical properties, the cell permeability, and the growth hormone that will finally promote plant

growth (Ampong et al., 2022). Humic acid will increase the permeability of membrane cells, respiration, photosynthesis, and elongation of roots (Akhmad et al., 2013) and improve the water holding capacity, soil structure, and microbial population (Shah et al., 2018) by increasing the soil biochemical activities; and increase soil nutrients availability (Yang et al., 2021).




Figure 2. Effect of HCB application on plant leaf area: (a) water spinach, (b) spinach, and (c) mustard green. D0 = control, D1 = 15 t HCB ha<sup>-1</sup>, D2 = 30 t HCB ha<sup>-1</sup>, D3 = 45 t HCB ha<sup>-1</sup>.

Humic acid also promotes several hormones, such as auxin and cytokinin, to enhance nutrient metabolism and photosynthesis (Nardi et al., 2021). Biochar was a habitat for beneficial microorganisms such as *Pseudomonas* sp. as P solubilizing and *Acetobacter* as N-fixing bacteria (Verdiana et al., 2016; Widowati et al., 2020). Biochar has a large number of micropores; hence, it functions as a habitat for microorganisms. A higher number of microorganisms in the soil will increase the availability of nutrients that plants in sufficient quantities easily absorb. The availability of sufficient nutrients in the soil will help the formation of the vegetative part of the plant. Also, the wider the leaf area, the more chlorophyll is formed, which increases photosynthesis. The process of photosynthesis will produce plant biomass in high quantities.

Biochar has the physical and chemical ability to supply several plant nutrients (Hidayat, 2015). Biochar can increase plant height and leaf area by increasing

nutrient uptake (Sa'adah and Islami, 2019). Combining biochar with humic acid effectively enhanced plant height and leaf area. Humic acid results from organic matter extraction with a functional group capable of converting unavailable nutrients to available ones. Humic acid can maintain soil moisture, prevent nutrient leaching, improve soil structure, and increase microbial activity (Sembiring, 2016).

The Least Significant Difference (LSD) test showed that there were no significant differences in fresh weight and dry weight in all three plant types (Table 1) due to the application of humic acid-coated biochar (HCB), but the humic acid-coated biochar (HCB) doses significantly affected the plant fresh and dry weight parameters (Table 1). The rate of HCB affected the yield parameters, even though there was no difference among doses of HCB on the plant weight. The application of 15 t HCB ha<sup>-1</sup> seemed to show the best dry and fresh weight (Table 1). This result is in line with the review of Rosa et al. (2014),

that humic acid doses significantly affected shoot growth, while the HCB source affected both the shoot and root growth. Humic acid will promote hormone production such as auxin, cytokinin, and metabolic enzymes; by improving nutrient uptake and leaf chlorophyll concentration, the roots, shoots, and yield will increase. The similar yields observed between the different doses of HCB happened because the application rates are determined by the environmental and soil conditions, source and composition of biochar and humic acid, and crop types. Rose et al. (2014) stated that combined humic acid and other substances such as mineral fertilizer and biochar would form nutrient-complex and subsequent crop uptake slowly, but their effect mostly depends on the humic acid source, application rate, and crop types.

Table 1. The effect of HCB on the fresh and dry weight of water spinach, spinach, and mustard green plants.

Treatments	Fresh weight (g)	Dry weight (g)
Water Spinach		
D0 (control)	48.98 a	11.55 a
D1 (15 t HCB ha <sup>-1</sup> )	79.67 b	15.59 a
D2 (30 t HCB ha <sup>-1</sup> )	82.33 b	17.27 a
D3 (45 t HCB ha <sup>-1</sup> )	64.67 ab	14.78 a
LSD 5%	24.71	6.72
Mustard Green		
D0 (control)	34.67 a	4.01 a
D1 (15 t HCB ha <sup>-1</sup> )	45.67 ab	5.76 b
D2 (30 t HCB ha <sup>-1</sup> )	53.00 b	6.92 c
D3 (45 t HCB ha <sup>-1</sup> )	51.33 b	5.33 b
LSD 5%	13.11	1.02
Spinach		
D0 (control)	31.67 a	6.53 a
D1 (15 t HCB ha <sup>-1</sup> )	49.33 b	11.30 b
D2 (30 t HCB ha <sup>-1</sup> )	46.33 b	13.82 b
D3 (45 t HCB ha <sup>-1</sup> )	46.00 b	8.66 ab
LSD 5%	13.09	4.46

Notes: Numbers followed by the same letters in the same column show that they are not significantly different. HCB = biochar coated with humic acid, LSD = least significant difference.

Humic acid is a nutrient supplement in the form of organic substances that have a complex molecular structure. It stimulates and activates biological and physiological processes for microbes in the soil (Ouni et al., 2014). According to Suwardi and Wijaya (2013), using humic acid will benefit agricultural soil because it can convert unavailable elements into available forms. In addition, it maintains moisture, resists the leaching of nutrients, and improves soil structure to promote microbial activity. Compared with biochar, it is suspected that humic acid has not played a role in the short term but will play an active role in the long term. Biochar is a nest for microorganisms that can increase soil microbial activity. As an amendment,

biochar is quite effective in improving physical properties in the short term, but new chemical properties will be seen in the long term (Lehman et al., 2011). Widowati et al. (2017) reported that combining biochar with NPK fertilizer increased organic carbon by 40%. Organic matter is the most important component of soil fertility, which plays an important role in maintaining quality, including soil ecology (Gajda et al., 2013; Benbi et al., 2015).

Biochar combined with other ingredients such as nitrogen and phosphorus is also effective in improving plant growth. Widowati et al. (2020) reported that biochar combined with organic fertilizers improved soil chemical properties. The use of HCB in this study is considered effective because of the role of humic acid, which provides nutrients. Ammal et al. (2020) proved that biochar could increase soil fertility due to the improved status of macro and micronutrients. Dume et al. (2016) also revealed that biochar could increase soil pH, cation exchange capacity, organic carbon, total nitrogen, exchangeable cations, and available phosphorus.

#### *Pb and Cu uptake by plants*

Trace metal contents of Pb and Cu in plants part (root and leaves) at the end of the experiment are presented in Figure 3. Generally, the concentrations of Pb and Cu were significantly higher in below-ground biomass (roots) than in above-ground biomass (leaves). The roots tend to accumulate more heavy metals than other plant parts. The plants have difficulties in translocating the heavy metals from roots to shoots. This condition will enhance root damage due to the high concentration of heavy metals. The Pb uptake amounts in spinach, water spinach, and mustard green roots were 0.023 g pot<sup>-1</sup>, 0.014 g pot<sup>-1</sup>, and 0.025 g pot<sup>-1</sup>, respectively. Pb contents in leaves of spinach, water spinach, and mustard green plants were 0.001 g pot<sup>-1</sup>, 0.001 g pot<sup>-1</sup>, and 0.001 g pot<sup>-1</sup>, respectively.

A similar result was also observed in the Cu content in the leaves, but the amount of Cu taken up by roots and leaves was relatively high compared with Pb. The amounts of Cu uptake in the roots of spinach, water spinach, and mustard green were 0.282 g pot<sup>-1</sup>, 0.443 g pot<sup>-1</sup>, and 0.143 g pot<sup>-1</sup> for spinach, water spinach, and mustard green, respectively. In comparison, the amounts of Cu uptake in the leaves were 0.020 g pot<sup>-1</sup>, 0.066 g pot<sup>-1</sup>, and 0.041 g pot<sup>-1</sup> for spinach, water spinach, and mustard green, respectively (Figure 3). The results showed that humic acid-coated biochar effectively reduced Pb and Cu uptake by spinach, water spinach, and mustard green by accumulating the Pb and Cu in their root parts.

Types of organic amendments and doses rate affect the heavy metal present in the shoots of a plant (Li et al. (2021). The highest Pb uptake (0.025 g pot<sup>-1</sup>) was observed in mustard green, and the lowest was in spinach at 0.0014 g pot<sup>-1</sup>. According to FAO (2014), the Pb level did not exceed the critical level (0.3 mg kg<sup>-1</sup>), while the Cu content exceeded the critical level

(10 mg kg<sup>-1</sup>). The highest Cu uptake in the plant roots was found in the water spinach (0.443 g pot<sup>-1</sup>), while on spinach and mustard green, Cu uptake was 0.282 g pot<sup>-1</sup> and 0.143 g pot<sup>-1</sup>. The difference in the uptake of Cu and Pb was because each plant had different morphological properties, although all plants were categorized as moderate heavy metal accumulators. The Bio Concentration Factor (BCF) value of each

plant for Pb was 0.29, 0.16, and 0.63 for spinach, water spinach, and mustard green, respectively; while the BCF value for Cu was 0.67, 1.01, and 0.98 for spinach, water spinach, and mustard green. This means that the BCF moderately transferred the heavy metals from the soil into the plant. BCF is the ratio of the concentration of heavy metals in roots and the concentration of heavy metals in the soil (Yoon et al., 2006).




Figure 3. Pb and Cu uptake by plants.

Results of this study also showed that mustard green was more effective than the other plants in stabilizing soil Pb, while the water spinach was more effective than the other plants in immobilizing Cu. The results of this study also showed that spinach, water spinach, and mustard green planted on heavy metal-contaminated soil could absorb relatively high heavy metals. Water spinach, mustard green, and spinach are remediation plants that are possible to absorb and reduce heavy metal content in the soil. However, most heavy metals accumulate in the roots, while the heavy metal content in leaves and stems is relatively low.

Heavy metals taken up by the plant will be transferred into the roots, stems, and leaves. Results of this study showed that the contents of heavy metals in roots, stems, and leaves were quite high (Figure 4). When the Cu is present in the soil and absorbed in large quantities, it will be dangerous to plants and humans because it is carcinogenic (Hardiana, 2009). Water spinach, mustard green, and spinach are accumulator plants that absorb heavy metals and transfer them to the roots, stems, and leaves. In general, plants have a root structure consisting of vascular tissue, including phloem, xylem, endodermis, and epidermis. The endodermis has a very large role in the deepest layer of the root because it has a Casparian band (Casparian strip). This controls the transfer of water and solutions, including those containing heavy metals. According to Alberto et al. (2013), plants can decontaminate heavy metals through roots transported to the top. Some heavy metals will be inhibited or prevented from spreading in this process. However, this depends on the number of contaminants available for uptake. Heavy metal accumulation is a determinant of its translocation to plant parts. Detoxification of heavy metals by plants can improve quality by increasing soil

physical, chemical, and biological properties. The absorbed heavy metals (non-essential) will enter plant cells through competition with other essential metals by taking over the binding site. This condition results in an interaction between the uptake of essential and non-essential metals that are toxic to plants. After entering the cell, heavy metals bind to phytochelatin and form a stable complex, which is then transported to the vacuole. This process will further detoxify (Komarek et al., 2013). The detoxification of heavy metals affects the soil, which will produce healthy plants.

#### Soil Pb and Cu reduction

Soil amendments such as biochar, humic acid, or biochar enriched with humic acid effectively decreased the availability of Cd, Cu, Pb, and As (Li et al., 2021). These amendments reduce the exchangeable fraction and increase the oxidation and residual fraction. Figure 4 shows spinach, water spinach, and mustard green plants reduced soil Pb between 40.04 and 87.28%, while Cu was between 8.63% and 40.23%. This is in line with the research result of other researchers that biochar can remediate Pb between 18.8% and 77.0% and Cu around 20% (Sizmur et al., 2011; Jiang et al., 2012). These results showed that applying humic acid-coated biochar is effective in immobilizing the heavy metals in polluted soils. The decline of Pb and Cu uptake in plant tissue was probably due to the immobilization process (physisorption, chemisorption, and precipitation) of the heavy metals by biochar and humic acid. Biochar effectively inhibited the activation of plants to metals and reduced the uptake of metals by plants. The immobilization of heavy metal by biochar will convert the heavy metal into more stable and less toxic forms.

But, the presence of humic acid promoted the formation of humic acid Pb and Cu water complex and prevented the formation of Pb and Cu hydroxide, thus increasing the availability and mobility of soil Pb and Cu.




Figure 4. Reduction of Pb and Cu contents in the soil studied.

The results showed that humic acid-coated biochar consistently reduced soil Pb and Cu concentrations. The reduction of soil Pb was greater than soil Cu. This means that humic acid-coated biochar is more efficient in stabilizing the Pb than the soil Cu. Biochar characteristics affect the immobilization mechanisms. Biochar characteristics, such as high surface area, specific group, and alkaline character, are essential in heavy metal sorption. Biochar characteristics such as surface heterogeneity, functional groups, and large surface area will stabilize heavy metals by absorbing their soil surface and enhancing the precipitation and surface sorption due to their microporous structure.

The bioavailability of metals can be reduced by humic acid due to their strong affinity and their ability to form a stable chelate with metal ions through their carboxylic groups and phenolic-OH. Metals have different affinities for binding humic acid, depending on their stability constant, and from this point of view, Pb (14.8) have a stronger affinity than Cu (13.3). This emphasizes that the binding order of metals to humic acid in contaminated soil is Pb>Zn~Cu>Cd (Gusiatin and Kulikowska, 2016). Rong et al. (2020) reported that Cu and Pb have two binding sites; thus, the complexing capacity order is Pb>Cu (Abate and Masini, 2001). Rong et al. (2020) found that a reduction of available Pb (20% to 39%) in the soil was higher than Cd (23% to 37%). The characteristics and behavior of Pb and Cu reflected the results obtained in this study, where less Pb was taken up by plants than Cu (Figure 4). This means that less Cu is immobilized by humic acid-coated biochar than Pb so that more Cu is in available form and can be taken up by plants.

## Conclusion

Applying humic acid-coated biochar (HCB) at a dose of 30 t ha<sup>-1</sup> increased the growth of water spinach,

spinach, and mustard green plants that were grown on soil containing the critical levels of Pb and Cu. The three plants took up Pb and Cu in the roots and leaves in the range of 1.85-8.02 mg kg<sup>-1</sup> and 45.60-59.82 mg kg<sup>-1</sup>. The application of HCB to the soil water spinach, mustard green, and spinach reduced soil Pb content from 40.04 to 87.28% and soil Cu from 8.63 to 40.23%.

## Acknowledgements

The authors thank the Ministry of Education, Culture, Research, and Technology of Indonesia for funding this study through the PTUPT scheme. The support of Tribhuwana Tunggal University and the University of Pembangunan Nasional Veteran Jawa Timur for providing facilities and technical support is highly appreciated.

## References

- Abate, G. and Masini, J.C. 2001. Acid-basic and complexation properties of a sedimentary humic acid. A study on the Barra Bonita reservoir of Tietê river, São Paulo State, Brazil. *Journal of the Brazilian Chemical Society* 12:109-116.
- Ahmad, I., Saquib, R.U., Qasim, M. and Saleem, M.A. 2013. Humic acid and cultivar effects on growth, yield, vase life, and corm characteristics of gladiolus. *Chilean Journal of Agricultural Research* 73:339-344, doi:10.4067/S0718-58392013000400002.
- Ahmad, M., Rajapaksha, A.U., Lim, J.E., Zhang, M., Bolan, N., Mohan, D., Vithanage, M., Lee, S.S. and Ok, Y.S. 2013. Biochar as a sorbent for contaminant management in soil and water: A review. *Chemosphere* 99:19-33, doi:10.1016/j.chemosphere.2013.10.071.
- Alberto, A.M.P. and Sigua G.C. 2013. Phytoremediation: a green technology to remove environmental pollutants. *American Journal of Climate Change* 2:71-86, doi:10.4236/ajcc.2013.21008.
- Ammal, A., Abunyewa, A.A. and Yeboah, E. 2020. Influence of integrated soil fertility management on the vegetative growth parameters of *Zea mays* in the Guinea savanna eco-zone of Ghana. *Journal of Agricultural Sciences* 65(2):187-197, doi:10.2298/JAS2002187A.
- Ampong, K., Thilakarathna, M.S. and Gorim, L.Y. 2022. Understanding the role of humic acids on crop performance and soil health. *Frontiers in Agronomy* 4:848621, doi:10.3389/fagro.2022.848621.
- Arancon, N.Q., Lee, S., Edwards, C.A. and Atiyeh, R. 2003. Effect of humic acids derived from cattle, food and paper-waste vermicompost on growth of green house plants. *Pedobiologia* 47:741-744.
- Atiyeh, R.M., Edwards, C.A., Metzger, J.D., Lee, S. and Arancon, N.Q. 2002. The influence of humic acids derived from earthworm-processed organic wastes on plant growth. *Bioresource Technology* 84:7-14.
- Benbi, D.K., Kiranvir, B., Amardeep, S.T. and Shivani, S. 2015. Sensitivity of labile soil organic carbon pools to long-term fertilizer, straw and manure management in rice-wheat system. *Pedosphere* 25:534-545, doi:10.1016/S1002-0160(15)30034-5.
- De Melo, B.A.G., Motta, F.L. and Santana, M.H.A. 2016. Humic acids: Structural properties and multiple functionalities for novel technological developments.



- Material Science and Engineering C* 62:967-974, doi:10.1016/j.msec.2015.12.001.
- Dume, B., Mosissa, T. and Nebiyu, A. 2016. Effect of biochar on soil properties and lead (Pb) availability in a military camp in South West Ethiopia. *African Journal of Environmental Science and Technology* 10(3):77-85, doi:10.5897/AJEST2015.2014.
- Ertani, A., Sambo, P., Nicoletto, C., Santagata, S., Schiavon, M. and Nardi, S. 2015. The use of organic biostimulants in hot pepper plants to help low input sustainable agriculture. *Chemical and Biological Technologies in Agriculture* 2, Article number: 11.
- FAO. 2014. International Food Standard. General Standard for Contaminants and Toxin in Food and Feed. Codex Stan. 193-1995).
- Gajda, A.M., Przewoka, B. and Gawryjok, K. 2013. Changes in soil quality associated with tillage system applied. *International Agrophysics* 27:133-141, doi:10.2478/v10247-012-0078-7.
- Gayathri, B., Srinivasamurthy, C.A., Vasanthi, B.G., Naveen, D.V., Prakash, N.B. and Bhaskar, S. 2019. Extraction and characterisation of humic acid from different organic wastes and its physicochemical properties. *International Journal of Chemical Studies* 7(6):769-775, doi:10.22271/chemi.2020.v8.i1k.8359.
- Gusiati, M.Z. and Kulikowska, D. 2016. Behaviours of heavy metals (Cd, Cu, Ni, Pb and Zn) in soil amended with composts. *Environmental Technology* 37(18):1-31, doi:10.1080/09593330.2016.1150348
- Haghighi, M., Kafi, M. and Fang, P. 2012. Photosynthetic activity and N metabolism of lettuce as affected by humic acid. *International Journal of Vegetable Science* 18:182-189.
- Hamzah, A., Hapsari, R.I. and Priyadarshini, R. 2017. The influence of rice husk and tobacco waste biochars on soil quality. *Journal of Degraded and Mining Lands Management* 5(1):1001-1007, doi:10.15243/jdmlm.2017.051.1007.
- Hamzah, A., Hapsari, R.I. and Wisnubroto, E.I. 2016. Phytoremediation of cadmium-contaminated agricultural land using indigenous plants. *International Journal of Environmental & Agricultural Research* 2(1):8-14.
- Hardiani, H. 2009. Potential of plants in accumulating Cu metal in soil contaminated by solid waste paper industry. *Buana Sains* 44(1):27-40 (in Indonesian).
- Hidayat, B. 2015. Remediation of contaminated soil by using biochar. *Jurnal Pertanian Tropik* 2(1):51-61, doi:10.32734/jpt.v2i1.2878 (in Indonesian).
- Jeffery, S., Verheijen, F.G.A., van der Velde, M. and Bastos, A.C. 2011. A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture Ecosystems & Environment* 144:175-187, doi:10.1016/j.agee.2011.08.015.
- Jiang, T.Y., Jiang J., Xu, R.K. and Li, Z. 2012. Adsorption of Pb (II) on variable charge soils amended with rice-straw derived biochar. *Chemosphere* 89(3):249-256, doi:10.1016/j.chemosphere.2012.04.028,
- Kalavrouziotis, I., Koukoulakis, P.H. and Kostakioti, E. 2011. Assessment of metal transfer factor under irrigation with treated municipal wastewater. *Fuel and Energy Abstracts* 103, doi:10.1016/j.agwat.2011.11.002.
- Komárek, M., Vaněk, A. and Ettler, V. 2013. Chemical stabilisation of metals and arsenic in contaminated soils using oxides-A review. *Journal Environmental Pollution* 172:9-22, doi:10.1016/j.envpol.2012.07.045.
- Lehmann, J., Rillig, M.C., Thies, J., Masiello, C.A., Hockaday, W.C. and Crowley, D. 2011. Biochar effects on soil biota, a review. *Soil Biology and Biochemistry* 43:1812-1836, doi:10.1016/j.soilbio.2011.04.022.
- Li, S., Sun, X., Li, S., Liu, Y., Ma, Q. and Zhou, W. 2021. Effects of amendments on the bioavailability, transformation and accumulation of heavy metals by pakchoi cabbage in a multi-element contaminated soil. *RSC Advances* 11:4395-4405, doi:10.1039/D0RA09358K.
- Liu, H., Zhang, T., Tong, Y., Zhu, Q., Huang, D. and Zeng, X. 2022. Effect of humic and calcareous substance amendments on the availability of cadmium in paddy soil and its accumulation in rice. *Ecotoxicology and Environmental Safety* 231:113186, doi:10.1016/j.ecoenv.2022.113186.
- Lu, K., Yang, X., Shen, J., Robinson, B., Huang, H., Liu, D., Bolan, N., Pei, J. and Wang, H. 2014. Effect of bamboo and rice straw biochars on the bioavailability of Cd, Cu, Pb and Zn to *Sedum plumbizincicola*. *Agriculture, Ecosystems & Environment* 191, doi:10.1016/j.agee.2014.04.010.
- Maurya, A., Kesharwani, L. and Mishra, M.K. 2018. Analysis of heavy metal in soil through atomic absorption spectroscopy for forensic consideration. *International Journal for Research in Applied Science & Engineering Technology* 6(VI):1188-1192.
- Nardi, S., Schiavon, M. and Francioso, O. 2021. Chemical structure and biological activity of humic substances define their role as plant growth promoters. *Molecules* 26:2256, doi:10.3390/molecules26082256.
- Unni, Y., Ghnaya, T., Montemurro, F., Abdellia, Ch. and Lakhdar, A. 2014. The role of humic substances in mitigating the harmful effects of soil salinity and improve plant productivity. *International Journal of Plant Production* 8(3):353-374.
- Park, S.H., Cho, H.J., Ryu, C. and Park, Y.-K. 2016. Removal of copper (II) in aqueous solution using pyrolytic biochars derived from red macroalga *Porphyra tenera*. *Journal of Industrial and Engineering Chemistry* 36:314-319.
- Qin, L., Wang, Q., Jiang, E. and Gao, Z. 2016. Study on biochar coated urea fertilizer with lignin adhesive modified by different solvents. *Transactions of the Chinese Society for Agricultural Machinery* 47(5), doi:10.6041/j.issn.1000-1298.2016.05.023.
- Rong, Q., Zhong, K., Huang, H., Li, C., Zhang, C. and Nong, X. 2020. Humic acid reduces the available cadmium, copper, lead, and zinc in soil and their uptake by tobacco. *Applied Sciences* 10:1077, doi:10.3390/app10031077.
- Rose, M.T., Patti, A.F., Little, K.R., Brown, A.L., Jackson, W.R. and Cavagnaro, T.R. 2014. A meta-analysis and review of plant-growth response to humic substances: Practical implications for agriculture. *Advances in Agronomy* 124:37-89, doi:10.1016/B978-0-12-800138-7.00002-4.
- Sa'adah, N. and Islami, T. 2019. Effect of application of various types of biochar and N fertilizer on soybean growth and yield (*Glycine max* L.). *Jurnal Produksi Tanaman* 7:2077-2083 (in Indonesian).
- Saputo, S., Yusuf, M. and Sabila, S. 2016. Distribution of Pb and Cu heavy metals in river estimate sediments



- Silugonggo, Batang District, Pati regency. *Journal Oseanografi* 5(2):180-190 (in Indonesian).
- Satpathy, D., Reddy, M. and Dhal, S. 2014. Risk assessment of heavy metals contamination in paddy soil, plants, and grains (*Oryza sativa* L.) at the east coast of India. *BioMed Research International* 2014:545473, doi:10.1155/2014/545473.
- Sembiring, J.V., Nelvia, N. and Yulia, A.E. 2016. Growth of oil palm seeds (*Elaeis guineensis* Jacq.) in the main nursery on Ultisol sub soil medium given humic acid and compost of oil palm empty bunches. *Jurnal Agroteknologi* 6(1):25-32, doi:10.24014/ja.v6i1.1373 (in Indonesian).
- Shah, Z.H., Rehman, H.M., Akhtar, T., Alsamadany, H., Hamooh, B.T., Mujtaba, T., Daur, I., Al Zahrani, Y., Alzahrani, H.A.S., Ali, S., Yang, S.H. and Chung, G. 2018. Humic substances: determining potential molecular regulatory processes in plants. *Frontier in Plant Science* 9:263, doi:10.3389/fpls.2018.00263.
- Shifaw, E. 2018. Review of heavy metals pollution in China in agricultural and urban soils. *Journal of Health and Pollution* 68(18):180607, doi:10.5696/2156-9614-8.18.180607.
- Singh, R., Singh, D.P., Kumar, N., Bhargava, S.K. and Barman, S.C. 2010. Accumulation and translocation of heavy metals in soil and plants from fly ash contaminated area. *Journal of Environmental Biology* 31(4):421-443.
- Sizmur, T., Wingate, J., Hutchings, T. and Hodson, M.E. 2011. *Lumbricus terrestris* L. does not impact on the remediation efficiency of compost and biochar amendments. *Pedobiologia* 54 (Supplement):S211-S216, doi:10.1016/j.pedobi.2011.08.008.
- Susilo, D.E.H. 2016. Identification of constant value of leaf shape for leaf area measurement using length cross width of the leaf of horticulture plant in peat soil. *Anterior Jurnal* 14(2):139-146 (in Indonesian).
- Suwardi and Wijaya, H. 2013. Increased production of food crops with humic acid active ingredients with zeolite as a carrier. *Jurnal Ilmu Pertanian Indonesia* 18 (2):79-84 (in Indonesian).
- Tsukahara, T., Ezaki, T., Moriguchi, J., Furuki, K.m Shimbo, S., Matsuda-Inoguchi, N. and Ikeda, M. 2003. Rice as the most influential source of cadmium intake among general Japanese population. *Science of the Total Environment* 305:41-51, doi:10.1016/S0048-9697(02)00475-8.
- Verdiana, M.A., Thamrin, H. and Sumarni, T. 2016. Effect of dose of rice husk biochar and NPK fertilizer on growth and yield of maize. *Jurnal Produksi Tanaman* 4(8):611-616 (in Indonesian).
- Wang, X., Sato, T., Xing, B. and Tao, S. 2005. Health risks of heavy metals to the general public in Tianjin, China via consumption of vegetables and fish. *Science of the Total Environment* 350(1-3):28-37.
- Widowati, Sutoyo, Iskandar, T. and Karamina, H. 2017. Characterisation of biochar combination with organic fertilizer: the effects on physical properties of some soil types. *Bioscience Research* 14(4):955-965.
- Widowati, Sutoyo, Karamina, H. and Wahyu, F. 2020. Soil amendment impact soil organic matter and physical properties of the three soil types after second corn cultivation. *AIMS Agriculture and Food* 5(1):150-168, doi:10.3934/agrfood.2020.1.150.
- Yang, F., Tang, C. and Antonietti, M. 2021. Natural and artificial humic substances to manage minerals, ions, water, and soil microorganisms. *Chemical Society Reviews* 50:6221-6239, doi:10.1039/D0CS01363C.
- Yoon, J., Cao, X. and Zhou, O. 2006. Accumulation of Pb, Cu and Zn in native plants growing on a contaminated Florida site. *Science of the Total Environment* 368(2-3):456-464, doi:10.1016/j.scitotenv.2006.01.016
- Zhang, M-K. and Ke, Z-X. 2004. Heavy metals, phosphorus and some other elements in urban soils of Hangzhou City, China. *Pedosphere* 14(2):177-185.
- Zhou, H., Meng, H., Zhao, L., Shen, Y., Hou, Y., Cheng, H. and Song, L. 2018. Effect of biochar and humic acid on the copper, lead, and cadmium passivation during composting. *Bioresource Technology* 258:279-286.