

**Research Article**

## **Response of selected chemical properties of extremely acidic soils on the application of limes, rice husk biochar and zeolite**

**Gina Aliya Sopha<sup>1\*</sup>, Catur Hermanto<sup>1</sup>, Huub Kerckhoffs<sup>2</sup>, Julian A Heyes<sup>3</sup>, James Hanly<sup>3</sup>**

<sup>1</sup> Research Center for Horticultural and Estate Crops, National Research and Innovation Agency (BRIN), Bogor, West Java, Indonesia

<sup>2</sup> Start-A-Fresh, Mount Manganui, New Zealand

<sup>3</sup> Massey University, Palmerston North 4472, New Zealand

\*corresponding author: ginasopha80@gmail.com, gina006@brin.go.id

### **Abstract**

#### *Article history:*

Received 10 July 2022

Accepted 14 September 2022

Published 1 October 2022

#### *Keywords:*

acidic soils  
aluminium toxicity  
ammeliorant  
liming

Extremely acidic soils have low pH, high concentration of exchangeable  $\text{Al}^{3+}$  and low cation exchange capacity (CEC) that cause severe growth for most plants. The study was conducted in the soil laboratory of the Indonesian Vegetable Research Institute, Lembang, from June to August 2019. A randomised complete block design with seventeen treatments, three replications, and three incubation times (3, 30 and 60 days) was deployed to assess the effect of rates of soil amendments, namely 5 to 30 t liming materials  $\text{ha}^{-1}$ , 5 to 20 t rice husk biochar  $\text{ha}^{-1}$ , and 5 to 20 t zeolite  $\text{ha}^{-1}$  on extremely acidic soils. The results showed that lime materials, i.e., lime, agriculture limestone, and hydrated lime had a similar effect on increasing soil pH and reducing exchangeable  $\text{Al}^{3+}$ . Calcium super seemed more effective in increasing soil pH and reducing exchangeable  $\text{Al}^{3+}$  than local lime due to the high CCE value. However, there was no significant response to the highest rice husk biochar and zeolite rate on soil pH and exchangeable  $\text{Al}^{3+}$ . Rice husk biochar increased the concentration of  $\text{K}^+$  and zeolite raised the concentration of  $\text{Na}^+$ . However, the effect was minimal. About 3 t lime  $\text{ha}^{-1}$  or 2.5 t calcium super  $\text{ha}^{-1}$  or equivalent to 1.5 to 2 times exchangeable  $\text{Al}^{3+}$  is required to obtain the soil pH target of 4.8, where the exchangeable  $\text{Al}^{3+}$  was less than  $0.5 \text{ cmol}_{(+) } \text{ kg}^{-1}$ .

**To cite this article:** Sopha, G.A., Hermanto, C., Kerckhoffs, H., Heyes, J.A. and Hanly, J. 2022. Response of selected chemical properties of extremely acidic soils on the application of limes, rice husk biochar and zeolite. *Journal of Degraded and Mining Lands Management* 10(1):4011-4017, doi:10.15243/jdmlm.2022.101.4011.

### **Introduction**

The majority of weathered tropical soils are leached, have high annual precipitation, and are acidic, with a soil pH of less than 5.2 (Juo and Franzluebbbers, 2003). About one-third of the tropical lands are very acidic and contain a high concentration of exchangeable aluminium (Al) that potentially harm crop growth (Cristancho et al., 2014).

Many upland shallot production areas in West Java are acidic, with soil pH below 5.5 (Sopha et al., 2019). Moreover, the natural process and human

disturbance accelerate acidifying process through precipitation, leaching, crop nutrient removal, soil organic matter, ion exchanges, chemical reactions and fertiliser application (Havlin, 2014). Soil amendments are used to improve soil chemical properties and reduce the negative effects of acidity. Liming is a common practice to increase soil pH and mitigate the adverse effect of soil acidity. However, the effectiveness of lime in improving the chemical properties of acidic soils varies, depending on the lime material characteristics, liming method and lime rate.

Lime application increases soil pH, and total base cations that increase base saturation (Fageria and Baligar, 2008; Cristancho et al., 2014). Moreover, the solubility of Al depends on the soil pH; the concentration of exchangeable  $Al^{3+}$  reduces when the soil pH increases (Moir and Moot, 2010). Application of 1 t lime  $ha^{-1}$  increased soil pH from 3.94 to 4.17 and reduced exchangeable  $Al^{3+}$  from 1.60 to 1.16  $cmol_{(+)}$   $kg^{-1}$  (Sopha et al., 2021). This Al value is still high, more than 0.5  $cmol_{(+)}$   $kg^{-1}$ , which might cause aluminium toxicity. Therefore, a higher rate of lime should be investigated further. Despite lime, biochar was reported to have a positive effect on soil chemical properties. A specific amount of elemental nutrients from biochar ash might be added due to biochar application and increased soil pH (Glaser et al., 2002). Zeolite also was reported give a positive impact on acidic soils. Application of 5 t zeolite  $ha^{-1}$  + 150 kg TSP  $ha^{-1}$  increased soil pH from 4.6 to 5.46 (Aainaa et al., 2018). However, the effect of biochar and zeolite

on soil properties could diversely depend on soil chemical properties and material characteristics.

This study aimed to identify the effect of lime and other soil amendments on soil pH, exchangeable  $Al^{3+}$ , base cations and CEC of extremely acidic soil from West Java, Indonesia.

## Materials and Methods

The research was conducted from June to August 2019 in the soil laboratory of the Indonesian Vegetable Research Institute, Lembang, West Java, Indonesia. Air-dried soil samples of 100 g were placed in a plastic cup, and all soil amendments were added. The soil amendments rates per ha were determined using a tillage depth of 20 cm and bulk density of 1020  $kg\ m^{-3}$ . The soil amendments treatments are presented in Table 1. The soil and soil amendments were mixed and wetted with deionised water to 70% of the field water holding capacity of the soil.

Table 1. The treatments of soil amendments.

| Treatments | Soil amendments                | Rate ( $t\ ha^{-1}$ ) | Rate ( $g\ 100\ g^{-1}\ soil$ ) |
|------------|--------------------------------|-----------------------|---------------------------------|
| 1          | Control                        | 0                     | 0                               |
| 2          | Local lime                     | 10                    | 0.5                             |
| 3          | Local lime                     | 20                    | 1.0                             |
| 4          | Local lime                     | 30                    | 1.5                             |
| 5          | Agricultural limestone         | 20                    | 1.0                             |
| 6          | Hydrated lime                  | 20                    | 1.0                             |
| 7          | Calcium Super                  | 5                     | 0.25                            |
| 8          | Calcium Super                  | 10                    | 0.5                             |
| 9          | Rice husk biochar              | 5                     | 0.25                            |
| 10         | Rice husk biochar              | 10                    | 0.5                             |
| 11         | Rice husk biochar              | 20                    | 1.0                             |
| 12         | Zeolite                        | 5                     | 0.25                            |
| 13         | Zeolite                        | 10                    | 0.5                             |
| 14         | Zeolite                        | 20                    | 1.0                             |
| 15         | Local lime + biochar           | 10 +10                | 0.5 + 0.5                       |
| 16         | Local lime + zeolite           | 10 +10                | 0.5 + 0.5                       |
| 17         | Local lime + biochar + zeolite | 10 + 5+ 5             | 0.5 + 0.25 + 0.25               |

Notes: Calcium carbonate equivalent (%) of local lime = 71%, agricultural limestone = 90%, hydrated lime = 109%, calcium super = 108%. pH of rice husk biochar = 7.07 and zeolite pH = 8.57. Calcium super contains 37% of CaO and 8% of MgO.

The soil was sampled at 0, 30, and 60 days from incubation and replicated three times for each incubation time. The soil was air-dried and passed through a 2 mm sieve before chemical characteristic analysis. The parameters are soil  $pH_{H_2O}$ , exchangeable  $Al^{3+}$ , base cations and CEC. Soil pH was measured using a Hanna pH meter (HI 2550) with a soil: water ratio of 1:5 (Miller and Kissel, 2010). Exchangeable  $Al^{3+}$  was extracted by KCl and was determined by the titrimetric method (Eviati and Sulaeman, 2009). The exchangeable base cations ( $Ca^{2+}$ ,  $Mg^{2+}$ ,  $K^+$ , and  $Na^+$ ) were extracted from the soil with  $NH_4^+$  using 1 M ammonium acetate pH 7 and the CEC was measured by spectrophotometry with  $\lambda = 636\ nm$  (Eviati and Sulaeman, 2009).

## Results and Discussion

### *pH, exchangeable $Al^{3+}$ , exchangeable $Ca^{2+}$ , and exchangeable $Mg^{2+}$*

The initial soil pH was 4.20, and exchangeable  $Al^{3+}$  was 1.68  $cmol_{(+)}$   $kg^{-1}$ . The base cation concentrations were low, being 2.60  $cmol_{(+)}$   $kg^{-1}$  of  $Ca^{2+}$ , 0.59  $cmol_{(+)}$   $kg^{-1}$  of  $Mg^{2+}$ , 0.60  $cmol_{(+)}$   $kg^{-1}$  of  $K^+$  and 0.04  $cmol_{(+)}$   $kg^{-1}$  of  $Na^+$ , which provided a 32% Base Saturation of the soil's CEC of 12  $cmol_{(+)}$   $kg^{-1}$ . All liming materials (lime, agricultural limestone, hydrated lime and calcium super) and the combination of lime and biochar or zeolite significantly increased soil pH in all incubation times compared to biochar and zeolite alone (Table 2). The control treatment soil, which did

not receive lime and other soil amendments, had an average soil pH between 4.12 to 4.33 over the 60 days duration of the incubation experiment. Three days after the incubation period, all treatments containing liming materials showed significant increases in soil pH. Application of 10, 20 and 30 t lime ha<sup>-1</sup> increased soil pH values to 5.64, 6.66 and 6.96 at 3 days after incubation, and the pH continued to increase over time, being 6.33, 7.20 and 7.73 at 60 days, respectively. The recommended optimum pH of vegetable crops is 6 to 6.5, which was achieved by 10 t local lime ha<sup>-1</sup> within 30 days of application.

All the liming materials succeeded in raising the soil pH above 5.0, and the exchangeable Al<sup>3+</sup> concentration downed to zero points (Table 2). The control treatment soil, which received no lime or other soil amendments, had an average exchangeable Al<sup>3+</sup> between 1.43 to 1.56 cmol<sub>(+)</sub> kg<sup>-1</sup> over the 60 days duration of the incubation experiment. The lowest rate of liming material was 5 t calcium super ha<sup>-1</sup> which decreased the exchangeable Al<sup>3+</sup> to 0 cmol<sub>(+)</sub> kg<sup>-1</sup>. This significant reduction in exchangeable Al<sup>3+</sup> was obtained with a lower rate of lime than in other studies, which might be due to the differences in soil buffering capacity and lime quality. For example, Andrade et al. (2002) reported that 8.8 t lime ha<sup>-1</sup> reduced exchangeable Al<sup>3+</sup> from 1.23 to 0.22 cmol<sub>(+)</sub> kg<sup>-1</sup> and Sopha et al. (2021) reported that the application of 1 t lime ha<sup>-1</sup> decreased the exchangeable Al<sup>3+</sup> from 0.27 to 0.04 cmol<sub>(+)</sub> kg<sup>-1</sup>.

All types and rates of liming materials increased the concentration of Ca<sup>2+</sup>, but this effect was not significant for rice husk biochar and zeolite treatments (Table 2). The control treatment soil, which received no soil amendments, had an exchangeable Ca<sup>2+</sup> between 2.40 and 3.62 cmol<sub>(+)</sub> kg<sup>-1</sup> over the 60 days duration of the incubation experiment. Three days after the incubation period, all treatments containing lime materials gave significant increases in exchangeable Ca<sup>2+</sup>. For the local lime treatments, the 10, 20 and 30 t lime ha<sup>-1</sup> achieved exchangeable Ca<sup>2+</sup> 14.95, 23.28 and 29.87 cmol<sub>(+)</sub> kg<sup>-1</sup> respectively. However, either all exchangeable Ca<sup>2+</sup> changes occurred within the first three days, with little or no change occurring between 30 and 60 days after the incubation period. On average, every 1 t lime ha<sup>-1</sup> increased the Ca<sup>2+</sup> concentration by 1.07 cmol<sub>(+)</sub> kg<sup>-1</sup>, while the calcium super increased the concentration of Ca<sup>2+</sup> by 1.09 cmol<sub>(+)</sub> kg<sup>-1</sup> after 60 days of incubation. A lower result was reported in sandy clay loam soil by Mosharraf et al. (2021) that every 1 t dolomite ha<sup>-1</sup> increased the concentration of exchangeable Ca<sup>2+</sup> by 0.27 cmol<sub>(+)</sub> kg<sup>-1</sup> after 60 days of incubation.

Calcium super, which was the only soil amendment to contain significant quantities of Mg<sup>2+</sup>, increased the concentration of exchangeable Mg<sup>2+</sup> in proportion to the amount added (Table 2). No other soil amendments changed the soil exchangeable Mg<sup>2+</sup> significantly. In addition, exchangeable Mg<sup>2+</sup> was

uninfected by incubation times, apart from a minor decline when biochar or zeolite was added in the presence of lime. The incorporation of calcium super enhanced the Mg<sup>2+</sup> linearly due to Mg concentration of calcium super. On average, every 1 t calcium super ha<sup>-1</sup> increased the concentration of Mg<sup>2+</sup> by 0.21 cmol<sub>(+)</sub> kg<sup>-1</sup>. The result was similar to Fungenzi et al. (2021), who reported a linear function for dolomite (22% MgO) addition on the concentration of exchangeable Mg<sup>2+</sup>.

#### *Exchangeable K<sup>+</sup> and Na<sup>2+</sup>, and CEC*

Biochar treatments raised the concentration of K<sup>+</sup> in incubated soils after three days of incubation time (Table 3). The increasing K concentration in the soil solution may come from cation substances from biochar. Rice husk biochar may contain about 175 mg kg<sup>-1</sup> of K (Milla et al., 2013), increasing K<sup>+</sup> concentration. A significant change was reported by Masulili et al. (2010) that the application of 10 t ha<sup>-1</sup> rice straw biochar increased K<sup>+</sup> ions by 0.31 cmol<sub>(+)</sub> kg<sup>-1</sup> from 0.21 to 0.51 cmol<sub>(+)</sub> kg<sup>-1</sup>. In this experiment, the application of 10 t ha<sup>-1</sup> rice husk biochar increased the concentration of K<sup>+</sup> by only about 0.08 cmol<sub>(+)</sub> kg<sup>-1</sup> from 0.67 to 0.75 cmol<sub>(+)</sub> kg<sup>-1</sup>. The difference result of K<sup>+</sup> changing might be due to the differences in pyrolysis temperature, heating rate, holding time and particle size that may affect the rice husk biochar characteristics (Ji-Lu, 2007). It is important to note that by 60 days of incubation, this small change was no longer present, and all incubated soils showed the same amount of exchangeable K<sup>+</sup>.

Zeolite significantly increased the exchangeable sodium (Na<sup>+</sup>) compared to other treatments (Table 3). If the exchangeable Na<sup>+</sup> was higher than 15% of cation exchange capacity (CEC), it might lead to serious soil physical problems, such as crusting that reduces water infiltration and inhibits plant growth (Laker and Nortje, 2019). In this experiment, the highest Na<sup>+</sup> saturation was 2.12% (Na<sup>+</sup> = 0.274 cmol<sub>(+)</sub> kg<sup>-1</sup>, CEC = 12.91 cmol<sub>(+)</sub> kg<sup>-1</sup>) which is much lower than the critical value. Therefore, it seems that zeolite application at this rate should not negatively impact plant growth. The increase in Na<sup>+</sup> due to zeolite content has previously been ascribed to Na<sub>2</sub>O in the zeolite, which dissolves when applied to soil (Wang et al., 2012). At the end of the incubation time (60 days after incubation), the cation exchange capacity (CEC) was improved significantly by the application of 20 t local and hydrated lime ha<sup>-1</sup>, 10 t calcium super and zeolite ha<sup>-1</sup>, 10 t local lime ha<sup>-1</sup> + 10 t biochar ha<sup>-1</sup> and 10 t local lime ha<sup>-1</sup> + 5 t biochar ha<sup>-1</sup> + 5 t zeolite ha<sup>-1</sup> but not with other treatments (Table 3). There was also no significant difference in CEC over time. It seems that in specific values, lime affected the CEC more significantly than biochar and zeolite. Application of lime increased CEC or negative charge commonly because of the detachment of H<sup>+</sup> from organic materials (Bolan et al., 2003).

Table 2. The effects of soil amendments and incubation time on soil pH, exchangeable Al<sup>3+</sup>, Ca<sup>2+</sup>, and Mg<sup>2+</sup>.

| Treatments<br>(*) | pH      |         |         | Al <sup>3+</sup> (cmol <sub>(+)</sub> kg <sup>-1</sup> ) |         |         | Ca <sup>2+</sup> (cmol <sub>(+)</sub> kg <sup>-1</sup> ) |          |          | Mg <sup>2+</sup> (cmol <sub>(+)</sub> kg <sup>-1</sup> ) |         |         |
|-------------------|---------|---------|---------|--|---------|---------|--|----------|----------|--|---------|---------|
|                   | 3 days  | 30 days | 60 days | 3 days   | 30 days | 60 days | 3 days   | 30 days  | 60 days  | 3 days   | 30 days | 60 days |
| 1                 | 4.16 d  | 4.33 g  | 4.12 e  | 1.43 b   | 1.56 b  | 1.51 a  | 2.40 e   | 3.56 f   | 3.62 f   | 0.54 c   | 0.59 cd | 0.57 c  |
| 2                 | 5.64 c  | 6.24 e  | 6.33 c  | 0.00 c   | 0.00 b  | 0.00 b  | 14.95 c  | 12.76 d  | 13.78 c  | 0.56 c   | 0.55 cd | 0.53 c  |
| 3                 | 6.66 ab | 7.27 c  | 7.20 b  | 0.00 c   | 0.00 b  | 0.00 b  | 23.38 b  | 17.44 c  | 21.14 b  | 0.57 c   | 0.56 cd | 0.54 c  |
| 4                 | 6.96 a  | 7.73 a  | 7.73 a  | 0.00 c   | 0.00 b  | 0.00 b  | 29.87 a  | 24.85 a  | 26.12 a  | 0.57 c   | 0.56 cd | 0.53 c  |
| 5                 | 6.82 a  | 7.39 bc | 7.41 ab | 0.00 c   | 0.00 b  | 0.00 b  | 23.56 b  | 19.55 bc | 24.63 ab | 0.52 c   | 0.51 d  | 0.51 c  |
| 6                 | 6.80 a  | 7.55 ab | 7.62 a  | 0.00 c   | 0.00 b  | 0.00 b  | 23.98 b  | 21.83 b  | 24.44 ab | 0.57 c   | 0.53 cd | 0.51 c  |
| 7                 | 5.64 c  | 5.64 f  | 5.49 d  | 0.00 c   | 0.00 b  | 0.00 b  | 8.31 de  | 7.80 e   | 8.16 de  | 1.75 b   | 1.91 b  | 1.77 b  |
| 8                 | 6.32 ab | 6.66 d  | 6.58 c  | 0.00 c   | 0.00 b  | 0.00 b  | 12.42 cd   | 11.32 d  | 12.08 cd | 2.27 a   | 2.75 a  | 2.67 a  |
| 9                 | 4.12 d  | 4.35 g  | 4.09 e  | 1.74 ab  | 1.56 a  | 1.34 a  | 2.22 e   | 3.88 f   | 3.69 ef  | 0.55 c   | 0.57 cd | 0.56 c  |
| 10                | 4.04 d  | 4.35 g  | 4.08 e  | 1.81 ab  | 1.62 a  | 1.53 a  | 2.09 e   | 3.70 f   | 3.67 e   | 0.57 c   | 0.59 cd | 0.59 c  |
| 11                | 4.13 d  | 4.35 g  | 4.13 e  | 1.69 ab  | 1.40 a  | 1.56 a  | 2.32 e   | 3.68 f   | 3.50 f   | 0.58 c   | 0.61 cd | 0.59 c  |
| 12                | 4.13 d  | 4.35 g  | 4.08 e  | 1.75 ab  | 1.51 a  | 1.46 a  | 2.21 e   | 3.76 f   | 3.70 ef  | 0.55 c   | 0.58 cd | 0.56 c  |
| 13                | 4.10 d  | 4.38 g  | 4.11 e  | 1.86 a   | 1.52 a  | 1.46 a  | 2.32 e   | 3.66 f   | 3.60 f   | 0.55 c   | 0.57 cd | 0.55 c  |
| 14                | 4.10 d  | 4.35 g  | 4.19 e  | 1.83 ab  | 1.50 a  | 1.51 a  | 2.38 e   | 3.54 f   | 4.02 ef  | 0.56 c   | 0.55 cd | 0.59 c  |
| 15                | 5.73 c  | 6.38 e  | 6.26 c  | 0.00 c   | 0.00 b  | 0.00 b  | 14.19 cd   | 12.90 d  | 13.98 c  | 0.58 c   | 0.56 cd | 0.53 c  |
| 16                | 5.74 c  | 6.38 e  | 6.32 c  | 0.00 c   | 0.00 b  | 0.00 b  | 14.33 cd   | 13.34 d  | 14.18 c  | 0.57 c   | 0.56 cd | 0.52 c  |
| 17                | 5.75 c  | 6.40 e  | 6.37 c  | 0.00 c   | 0.00 b  | 0.00 b  | 14.74 c  | 12.80 d  | 13.80 c  | 0.58 c   | 0.56 cd | 0.51 c  |
| <i>p</i> -value   | <0.01   | <0.01   | <0.01   | <0.01  | <0.01   | <0.01   | <0.01  | <0.01    | <0.01    | <0.01  | <0.01   | <0.01   |
| CV (%)            | 8.6     | 1.1     | 2.0     | 19.5   | 16.7    | 15.8    | 13.9   | 8.8      | 12.8     | 5.5  | 3.5     | 7.0     |

Notes: Treatments (\*) see Table 1. Tukey's method for means comparison; means presenting the same small letters in the same column and the same capital letters in the same row, are not significantly different at  $\alpha=5\%$ .

Table 3. The effect of soil amendments and incubation time on exchangeable K<sup>+</sup>, Na<sup>2+</sup>, and CEC.

| Treatments<br>(*) | K <sup>+</sup> (cmol <sub>(+)</sub> kg <sup>-1</sup> ) |            |         | Na <sup>+</sup> (cmol <sub>(+)</sub> kg <sup>-1</sup> ) |         |         | CEC(cmol <sub>(+)</sub> kg <sup>-1</sup> ) |           |            |
|-------------------|--|------------|---------|---|---------|---------|--|-----------|------------|
|                   | 3 days   | 30 days    | 60 days | 3 days  | 30 days | 60 days | 3 days                                     | 30 days   | 60 days    |
| 1                 | 0.68 ef  | 0.74 abcd  | 0.67 a  | 0.06 cd   | 0.05 c  | 0.06 c  | 11.9 ns                                    | 11.4 d    | 11.5 e     |
| 2                 | 0.67 f   | 0.71 bcde  | 0.62 ab | 0.05 d  | 0.05 c  | 0.07 c  | 11.2                                       | 11.8 cd   | 12.4 abcde |
| 3                 | 0.67 f   | 0.71 bcde  | 0.64 ab | 0.05 d  | 0.05 c  | 0.04 c  | 13.1                                       | 13.2 ab   | 13.2 abc   |
| 4                 | 0.66 f   | 0.71 bcde  | 0.59 ab | 0.05 d  | 0.04 c  | 0.07 c  | 12.9                                       | 12.8 abcd | 12.7 abcde |
| 5                 | 0.68 ef  | 0.73 bcde  | 0.56 ab | 0.09 cd   | 0.05 c  | 0.05 c  | 12.6                                       | 12.5 abcd | 12.4 abcde |
| 6                 | 0.69 def   | 0.73 abcde | 0.60 ab | 0.06 cd   | 0.06 bc | 0.06 c  | 14.1                                       | 13.4 a    | 12.7 abcd  |
| 7                 | 0.68 ef  | 0.73 bcde  | 0.58 ab | 0.06 cd   | 0.05 c  | 0.05 c  | 13.1                                       | 12.8 abcd | 12.2 abcde |
| 8                 | 0.67 f   | 0.70 bcde  | 0.46 b  | 0.06 cd   | 0.04 c  | 0.04 c  | 12.2                                       | 12.8 abcd | 13.5 ab    |
| 9                 | 0.75 a-d   | 0.79 abc   | 0.72 a  | 0.06 cd   | 0.06 bc | 0.07 c  | 11.9                                       | 11.7 cd   | 11.5 de    |
| 10                | 0.78 ab  | 0.79 ab    | 0.71 a  | 0.07 cd   | 0.06 bc | 0.06 c  | 12.1                                       | 11.9 bcd  | 11.6 cde   |
| 11                | 0.80 a   | 0.84 a     | 0.75 a  | 0.07 cd   | 0.05 c  | 0.06 c  | 12.0                                       | 11.8 bcd  | 11.7 cde   |
| 12                | 0.67 f   | 0.69 cde   | 0.60 ab | 0.12 cd   | 0.09 bc | 0.13 bc | 12.8                                       | 12.5 abcd | 12.2 abcde |
| 13                | 0.71 cdef  | 0.66 de    | 0.62 ab | 0.11 cd   | 0.14 bc | 0.18 b  | 12.1                                       | 12.4 abcd | 12.7 abcd  |
| 14                | 0.69 cdef  | 0.64 e     | 0.63 ab | 0.30 a  | 0.22 a  | 0.31 a  | 12.2                                       | 11.8 bcd  | 11.5 de    |
| 15                | 0.75 abc   | 0.76 a-d   | 0.65 ab | 0.07 cd   | 0.07 bc | 0.05 c  | 12.7                                       | 12.9 abc  | 13.1 abcd  |
| 16                | 0.68 ef  | 0.66 de    | 0.59 ab | 0.21 b  | 0.11 bc | 0.18 b  | 12.6                                       | 12.7 abcd | 12.7 abcde |
| 17                | 0.73 b-e   | 0.73 abc   | 0.62 ab | 0.13 c  | 0.09 bc | 0.14 b  | 12.7                                       | 13.3 a    | 13.9 a     |
| <i>p</i> -value   | <0.01  | <0.01      | <0.01   | <0.01   | <0.01   | <0.01   | 0.39                                       | 0.04      | 0.01       |
| CV (%)            | 2.6  | 4.8        | 9.6     | 24.2  | 34.4    | 36.0    | 8.80                                       | 5.84      | 6.93       |

Notes: Treatments (\*) see Table 1. Tukey's method for means comparison; means presenting the same small letters in the same column and same capital letters in the same row, are not significantly different at  $\alpha=5\%$ .

The effect was more significant when the soils contain high clay minerals with pH-dependent surface charge (Cherian and Arnepalli, 2015). Exchangeable  $\text{Al}^{3+}$  was quantified when the soil pH was below 5.42 (Figure 1). No data of exchangeable  $\text{Al}^{3+}$  were found in the soil pH range of 4.45 and 5.42. However, from the

trendline, it was determined that when the pH was above 4.8, the concentration of exchangeable  $\text{Al}^{3+}$  was lower than  $0.5 \text{ cmol(+) kg}^{-1}$ . The point is the threshold level when the adverse effect of  $\text{Al}^{3+}$  on crop growth is negligible (Sopha et al., 2021). However, this finding should be investigated further.




Figure 1. The scatter points between  $\text{pH}_{\text{H}_2\text{O}}$  and exchangeable  $\text{Al}^{3+}$  concentration,  $n = 153$ .

## Conclusion

Lime materials, i.e. lime, agricultural limestone and hydrated lime, had a similar effect on increasing soil pH and reducing the exchangeable  $\text{Al}^{3+}$  in extremely acidic soil. In contrast, calcium super seems more effective in increasing soil pH and reducing exchangeable  $\text{Al}^{3+}$  than local lime. To reach the soil pH target, 4.8 (exchangeable  $\text{Al}^{3+}$  of  $<0.5$ ), about 3 t lime  $\text{ha}^{-1}$  or 2.5 t calcium super  $\text{ha}^{-1}$  or equivalent to 1.5 to 2 times exchangeable  $\text{Al}^{3+}$  is required.

## Acknowledgements

The authors thank the Indonesian Vegetable Research Institute for providing laboratory equipment. Thanks are also forwarded to the Indonesian Agency of Agriculture Research and Development for the scholarship and research fund. A part of this article was presented at the 2<sup>nd</sup> International Conference on Environment, Socio-Economic, and Health Impacts of Degraded and Mining Lands, 30-31st August 2022, Universitas Brawijaya, Malang, Indonesia.

## References

- Aainaa, H.N., Ahmed, O.H. and Ab Majid, N.M. 2018. Effects of clinoptilolite zeolite on phosphorus dynamics and yield of *Zea mays* L. cultivated on an acid soil. *PLoS One* 13(9):2-19, doi:10.1371/journal.pone.0204401.
- Andrade, D., Murphy, P. and Giller, K. 2002. Effects of liming and legume/cereal cropping on populations of indigenous rhizobia in an acid Brazilian Oxisol. *Soil Biology and Biochemistry* 34(4):477-485.

- Bolan, N., Adriano, D. and Curtin, D. 2003. Soil acidification and liming interactions with nutrient and heavy metal transformation and bioavailability. *Advances in Agronomy* 78:215-272, doi:10.1016/S0065-2113(02)78006-1.
- Cherian, C. and Arnepalli, D.N. 2015. A critical appraisal of the role of clay mineralogy in lime stabilisation. *International Journal of Geosynthetics and Ground Engineering* 1:8, doi:10.1007/s40891-015-0009-3.
- Cristancho, R., Hanafi, M., Syed Omar, S. and Rafii, M. 2014. Aluminum speciation of amended acid tropical soil and its effects on plant root growth. *Journal of Plant Nutrition* 37(6):811-827, doi:10.1080/01904167.2014.881856.
- Eviati and Sulaeman. 2009. *Chemical Analysis of Soil, Plants, Water and Fertilisers* (Technical Guide Edition 2). Soil Research Institute, Bogor (in Indonesian).
- Fageria, N. and Baligar, V.C. 2008. Ameliorating soil acidity of tropical Oxisols by liming for sustainable crop production. In: Sparks, D. (ed.), *Advances in Agronomy* Vol. 99. Academic Press is an imprint of Elsevier, pp 345-399, doi:10.1016/S0065-2113(08)00407-0.
- Glaser, B., Lehmann, J. and Zech, W. 2002. Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal – a review. *Biology and Fertility of Soils* 35(4):219-230, doi:10.1007/s00374-002-0466-4.
- Havlin, J.L., Tisdale, S.L., Nelson, W.L. and Beaton, J.D. 2014. *Soil Fertility and Fertilisers: An Introduction to Nutrient Management*. Pearson.
- Ji-Lu, Z. 2007. Bio-oil from fast pyrolysis of rice husk: Yields and related properties and improvement of the pyrolysis system. *Journal of Analytical and Applied Pyrolysis* 80(1):30-35.

- Juo, A.S. and Franzluebbers, K. 2003. *Tropical Soils: Properties and Management for Sustainable Agriculture*. Oxford University Press., doi:10.1111/j.1365-2389.2004.0635c.x.
- Laker, M.C. and Nortjé, G.P. 2019. Review of existing knowledge on soil crusting in South Africa. In: Sparks, D.L. (Ed.), *Advances in Agronomy* Vol. 155. Academic Press. pp. 189-242, doi:0.1016/bs.agron.2019.01.002.
- Masulili, A., Utomo, W.H. and Syekhfani, M.S. 2010. Rice husk biochar for rice-based cropping system in acid soil. 1. The characteristics of rice husk biochar and its influence on the properties of acid sulfate soils and rice growth in West Kalimantan, Indonesia. *Journal of Agricultural Science* 2(1):39-47, doi: 10.5539/jas.v2n1p39
- Milla, O.V., Rivera, E.B., Huang, W.J., Chien, C.C. and Wang, Y.M. 2013. Agronomic properties and characterisation of rice husk and wood biochars and their effect on the growth of water spinach in a field test. *Journal of Soil Science and Plant Nutrition* 13:251-266.
- Miller, R.O. and Kissel, D.E. 2010. Comparison of soil pH methods on soils of North America. *Soil Science Society of America Journal* 74(1):310-316.
- Moir, J. L. and Moot, D.J. 2010. *Soil pH, Exchangeable Aluminium and Lucerne Yield Responses to Lime in a South Island High Country Soil*. The New Zealand Grassland Association 72.
- Mosharrof, M., Uddin, M., Jusop, S., Sulaiman, M.F., Shamsuzzaman, S. and Haque, A.N.A. 2021. Changes in acidic soil chemical properties and carbon dioxide emission due to biochar and lime treatments. *Agriculture* 11(3):219, doi:10.3390/agriculture11030219
- Sopha, G.A., Hermanto, C., Hanly, J., Heyes, J.A. and Kerckhoffs, H. 2019. A Survey of Soil Fertility and Shallot (*Allium cepa* Aggregatum Group) Yields in West Java, Indonesia. *Proceedings of The Emerging Challenges and Opportunities in Horticulture Supporting Sustainable Development Goals ISH 2018*. November 27-30, 2018, Kuta, Bali, Indonesia. p 341-346, doi: 10.26352/CY27\_ISH-2018.
- Sopha, G.A., Hermanto, C., Hanly, J., Heyes, J.A. and Kerckhoffs, H. 2021. Influence of lime and phosphorus fertiliser on shallot growth and bulb yield in strongly acid soils on West Java, Indonesia. *Acta Horticulturae* 1312:315-322, doi:10.17660/ActaHortic.2021.1312.46.
- Wang, X., Ozdemir, O., Hampton, M.A., Nguyen, A. V. and Do, D.D. 2012. The effect of zeolite treatment by acids on sodium adsorption ratio of coal seam gas water. *Water Research* 46(16): 5247-5254.