

Research Article

Hydroseeding application using pioneer local plant seeds for coal postmining soil in Tanah Laut Regency, South Kalimantan

M.F. Anshari^{1*}, E. Boedianto², A.A.R. Fernandes³, E. Arisoesilaningsih¹

¹ Biology Master Program, Faculty of Mathematics and Natural Sciences, Brawijaya University, Jl. Veteran, Malang, East Java, Indonesia

² PT Amanah Anugerah Adi Mulia Site Kintap, Riam Adungan Village, Kintap District, Tanah Laut Regency, South Kalimantan

³ Department of Statistics, Faculty of Mathematics and Natural Sciences, Brawijaya University, Jl. Veteran, Malang, East Java, Indonesia

*corresponding author: delansh@gmail.com

Received 28 April 2018, Accepted 19 May 2018

Abstract: The research aimed to determine pioneer local plant seeds germination and growth on coal postmining soil, and to determine optimal mulch for producing great productivity. The research was conducted in two applications in a coal post mining area, Tanah Laut, South Kalimantan. The used seeds were 5 species of legumes, 3 species of grasses, and 3 species of sedges. Hydroseeding was applied by mixing seeds in mulch and spreading it on soil surface of 400 cm² plots (1st application) and 10,000 cm² plots (2nd application). There were five treatments in the 1st application and repeated four times, while in the 2nd, there were six treatments with three times repetitions. Germination rate, seedling density, plant coverage, height/length, and leaf length were observed every three days (1st application). On 60 days after sowing, root length and plant biomass (the 1st), and plant coverage, height and productivity (the 2nd) were measured. The results showed that hydroseeding mulches could be media for pioneer local plant seeds germination and seedling growth especially for Leguminosae. While the buffalo dung could be hydroseeding mulch for Poaceae growth. Although root/shoot ratio of Cyperaceae was higher than other families, Leguminosae root developed longer as shoot growth and well adapted on postmining soil. The hydroseeding mulch composed by 720 seeds per plot and applied 4 cm in depth was optimal based on seed germination percentage. Whereas mulch with 0.2 cm depth in 2nd application showed a higher effectivity for plant growth and productivity.

Keyword: *hydroseeding, local plants, mining soil, revegetation*

To cite this article: Anshari, M.F., Boedianto, E., Fernandes, A.A.R. and Arisoesilaningsih, E. 2018. Hydroseeding application using pioneer local plant seeds for coal postmining soil in Tanah Laut Regency, South Kalimantan. *J. Degrade. Min. Land Manage.* 5(4): 1335-1345, DOI: 10.15243/jdmlm. 2018.054.1335.

Introduction

Mining has become the biggest contributor to the economic development of Indonesia for about last 30 years (Manaf, 2009). The main area of coal deposits in Indonesia are Northern Sumatra (54%), Eastern Kalimantan (28%), Southern Kalimantan (10%), Riau (2%), and Central Kalimantan (1.4%) (Ministry of Energy and Mineral Resources, 2008). Based on data of the General Directorate of Fiscal Balance at the Ministry of Finance of the Republic of Indonesia

(2012), the mining sector in Southern Kalimantan contributes revenue of 24.42% of the total Gross Domestic Product (GDP) in 2011, which reached IDR 68,234 billion. Coal mining produces 141,809,435 million tonnes from eight producer districts, namely Tanah Laut, Kota Baru, Banjar, Tapin, Hulu Sungai Selatan, Tabalong, Tanah Bumbu, and Balangan.

Generally, the technique of coal mining in Kalimantan uses open pit mining techniques with back filling methods (Darmawan and Irawan,

2009; Subandrio et al., 2009). The negative impact of mining is the loss of soil cover because topsoil and subsoil are reversed and moved, while the main material of rocks appears on surface. The oxidation process of sulfide minerals such as pyrite will release sulfuric acids that have negative impacts for decreasing drastically soil pH and affecting the balance of nutrients in the soil (Rochani and Damayanti, 1997). Topsoil displacement also causes loss of soil organic matter, while nutrients will be leached when rain falls. Nutrient and organic matter poor soil has low capacity for plant growth, fertilizer application and water holding (Djajakirana, 2001).

Post-mining land reclamation and revegetation are held to overcome negative impacts of mining activities because of the open pit mining techniques. It aims to improve the disturbed land use after mining operations, as well as to accelerate plant coverage and its ecological service as intended (Decree of the Minister of Energy and Mineral Resources No. 1211.K/008/M.PE/1995; Decree of the Minister of Forestry and Plantation No. 146/Kpts-II/1999). Among revegetation methods, hydroseeding becomes promising. Hydroseeding is a planting process using a mixture of seeds and mulch that mixed together and sprayed on the land that has been prepared. Unstable soil surface due to the displacement of vegetation, native soil, and debris on coal mining was successfully revegetated with hydroseeding using grasses, legumes and fertilizer nutrients (Zipper et al., 2012).

Revegetation as a part of diversity restoration program increases the accumulation of carbon and structures by involving direct carbon input through mulching and transplanting. These efforts are particularly applied by using fast growing ground cover plants that can stabilize the soil surface (Walker and del Moral, 2008). However the local plants as seed sources in post mining land revegetation is greatly selected due to its adaptation to local biotic and abiotic factors (Oliveira et al., 2013). Meanwhile, in Tanah Laut Regency, a herd of buffalo that visits and grazes in post-mining land left dung that is essentially an example of a natural hydroseeding. In a few weeks, various grasses grow naturally from the decomposed dung. Therefore, it is interesting to use the buffalo dung hydroseeding model and imitate it for an artificial hydroseeding using pioneer local plant seeds using different mulches composition in coal post-mining soil. The aims of this study were to evaluate the hydroseeding applications using local plants seeds and some mulches composition, determining its germination and growth in coal post-mining soil, and selecting

the optimal mulch likes buffalo dung performance.

Materials and Methods

A field research was conducted in two steps of application. The first one was held from January to April 2015 in the rainy season and the second step was held from February to May 2017, in PT Amanah Anugerah Adi Mulia Site Kintap. It is located in Riam Adungan Village, Kintap District, Tanah Laut Regency, South Kalimantan, with four points of coordinate boundaries of Production Operation Mining Permits (IUP.OP) and approximately 33 km from Southern of Trans South Kalimantan Road (Figure 1). The mining area is around 599.78 ha, and bordered on the North by IUP of PT WTB, on the West by IUP of KUD TJM, on the South by IUP of PT Ananta, and on the East by the areas of limited production forest. The nearest villages are Riam Adungan and Salaman within \pm 6-10 km. The annual rainfall is 1,771.5 mm.

The collected seeds that were dried consisted of five species of Leguminosae (*Cajanus cajan* (L.) Millsp., *Crotalaria pallida* Aiton., *Desmodium triflorum* (L.) DC., *Indigofera spicata* Auct., *Sesbania grandiflora* (L.) Pers.), three species of Poaceae (*Eleusine indica* (L.) Gaertn., *Sporobolus indicus* (L.) R. Br., *Paspalum conjugatum* PJ Bergius), and three species of Cyperaceae (*Cyperus brevifolius* (Rottb.) Endl. ex Hassk., *Cyperus odoratus* L., *Kyllingia monocephala* Rottb.). These pioneers were found growing wild in the post coal-mining area of PT Amanah. The seeds were soaked with water for 24 hours. The sinking seeds were then treated with Gibberellic Acid (GA3) for 1 hour and mixed with mulch. Number of seeds used for 360 seeds density was composed by 60 seeds of Leguminosae, 150 seeds of Cyperaceae, and 150 seeds of Poaceae. Number of seeds used for 720 seeds density was consisted by 120 seeds of Leguminosae, 300 seeds of Cyperaceae, and 300 seeds of Poaceae.

According to the results of preliminary studies, mulch of the 1st application was made from a mixture of wood fibers, water, local microorganisms (mol), adhesive, compost, charcoal, and ash. Mulch was then added with seeds of all plant species proportionally with two different densities, 360 seeds (B₁) or 720 seeds (B₂) per plot. After mixing, the mulch was spread over 20 cm x 20 cm on the post-mining soil plots with two levels of depth, which were 4 cm (M₁) and 6 cm (M₂), and a control (fresh buffalo dung with 2 cm depth). All treatments were repeated

four times, while watering and observation were held every three days for 2 months. For the 2nd application, mulch (M₃) was composed by tackifier agent, water, compost, rice husk,

sawdust, and urea. Mulch was mixed with plant seeds proportionally and stirred up until forming a slurry mixture, and then it was spread over 100 cm x 100 cm plots. It was repeated three times.




Figure 1. Hydroseeding application site in coal postmining soil at PT Amanah, Tanah Laut Regency, South Kalimantan

In the 1st application we observed seeds germination time in every planting medium. The number of germinated seeds was divided by the sown seeds number to determine the seed germination rate. The growth variables of seedling included plant height, leaf length, root length, biomass (shoot and root), and productivity. Vegetation analysis was conducted to determine density and coverage of Leguminosae, Cyperaceae, and Poaceae. On 60 days after sowing, all seedlings were harvested, we determined root length, biomass and productivity. While in 2nd application we observed plant coverage, height, and productivity.

Germination and seed growth data such as height, number of leaves, root length, density and coverage, biomass (shoot and root), and productivity were compiled using Microsoft Excel program. Furthermore, data of density, coverage, root/shoot ratio, root nodules, and productivity, were univariately statistical analyzed. While data of germination and growth variations on some planting media were analyzed multivariately by biplot and cluster analysis using PAST software (Hammer et al., 2001).

Results and Discussion

Seeds germination of some pioneer local plants on mulches

The prepared hydroseeding mulches could be served as seeds germination media especially

Leguminosae seeds. While the fresh buffalo dung mulch as control, could be Poaceae seed germination medium (Figure 2). Mulch depth and seed density affected seedling number. A thinner mulch (M₁) produced more seedling number than M₂. Thick mulch and sun were observed as limiting factors of grass seed germination. In addition, higher density seeds mulches (B₂) also produced more seedling number than B₁ mulches (Table 1). Nevertheless, the seedling number in this experiment was still very small, ranged 1-5%. However in the greenhouse experiment, the most Leguminosae seeds produced 40% seedling on 12 days after sowing (Baiti and Arisoesilansih, 2015). Some possible factors produced this low germination rate of Leguminosae seed in post-mining land were fungal attack, low viability of seeds, and seed predation by ants (Anshari, 2015). Fungal attack occurred if the medium was very acidic and moist. In addition, immature seeds germination stimulated the fungi growth (Utomo, 2006). According to Abella et al. (2009), some variables influencing seed germination success were selection of species, genetic stock, and rate of germination, given treatment, environmental conditions, germination time, and rainfall. To overcome this low seeds germination, increasing number and better selection of seeds could be conducted. Soaking seeds in hot water with certain temperature and duration is another method to raise seed germination rate, particularly for Leguminosae seeds (Kak et al., 2007; Naikawadi, 2016).




Figure 2. Seed germination of some pioneer local plants at 15 days after sowing on mulches

Note: B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; Control: Fresh buffalo dung with 2 cm depth. Scale represents 2 cm.

Table 1. Cumulative seedling number of three pioneer local plant families grown on mulches until 15 days after sowing

No	Families	Total Seedling (individual)				
		B ₁ M ₁	B ₁ M ₂	B ₂ M ₁	B ₂ M ₂	Control
1	Leguminosae/Broadleaf plant	3 (5)	2 (3)	6 (5)	6 (5)	1
2	Cyperaceae	3 (2)	4 (3)	12 (4)	6 (2)	2
3	Poaceae	3 (2)	2 (1)	12 (4)	4 (1)	99

Note: Numbers in parenthesis indicate relative seedling number (%), compared to the sown seeds; B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; Control: Fresh buffalo dung with 2 cm depth.

Seedling growth of some local plant families on mulches


Density of Cyperaceae seedling until 42 days after sowing tended to increase irregularly in each observation. The highest density was recorded on B₂M₁ treatment and reached 18 individuals/400 cm², followed by B₂M₂ treatment reached 13 individuals/400 cm². Moreover in Leguminosae/broadleaf plants, seedling density until 42 days after sowing tended to fluctuate due to herbivore attack or seedling death. The highest seedling density was observed in B₂M₂ treatment and reached 9 individuals/400 cm², while in B₂M₁ treatment it reached 7 individuals/400 cm² (Figure 3A). Differently with two previous families, Poaceae seedling density until 42 days after sowing tended to increase, with the highest density in control plots that reached 355 individuals/400 cm². While in all treatments, Poaceae density was very low compared to control, respectively 24 individuals/400 cm² and 10 individuals/400 cm² in B₂M₁ and B₂M₂, (Figure 3B). These results revealed that seedling

density in the mulch with 720 seeds density (B₂) was significantly higher than seedling density in the mulch with 360 seeds density (B₁). Seed quality and dormancy, inhibitors presence, and predation might serve as limiting factors (Anshari, 2015). However, the mulch in control was more suitable for Poaceae growth medium. It is assumed that it closely related with buffalo's diet. Lancelotti and Madella (2012) showed that buffalo dung was composed by grass leaf/culm and grass inflorescence/seeds due to commonly the buffaloes graze in the pasture lands. Poaceae is the most important fodder for buffalo (Harun et al., 2017). From 53 species of Poaceae, there were 27 species of buffalo fodder in Central Punjab. Additionally, Zereen et al. (2013) stated that Poaceae played a significant fodder because almost all of 51 species of Poaceae in Central Punjab, were used as fodder for the animals, including buffalo.


Coverage of some local plant families increased until 60 days after sowing. The highest Leguminosae coverage was seen in B₂M₂ treatment and reached 36.25% (Figure 4A). While

in Cyperaceae and Poaceae, the highest coverage found in M₃ treatment, respectively 13.84% and 37.74%. In the controls, Poaceae dominated with highest coverage and reached 78.75% (Figure 4B-C and Figure 5). This result was still less than those reported by Fan et al. (2013). They showed that Poaceae coverage on 60 days after application reached more than 90%. The control mulch could be similar as compost blanket as reported by Faucete et al. (2006). They revealed that blanket compost with 3.75 cm depth provided vegetation

coverage 2.75 times greater in three months compared to hydroseeding treatment. This greater vegetation coverage on the compost treatment was likely due to their ability to maintain moisture. Then the lower coverage on hydroseeding was possibly caused by drained seeds during rainfall and or greater evaporation after hydroseeding treatment. Therefore erosion control could be provided by combining compost and a thicker hydroseeding application.



A



B

Figure 3. Seedling density of pioneer local Cyperaceae, Leguminosae/broadleaf plant, and Poaceae until 42 days after sowing on mulches. Plot size is 400 cm².

Note: A: Cyperaceae and Leguminosae; B: Poaceae; C: Cyperaceae; L/DL: Leguminosae/broadleaf plant; P: Poaceae; B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; Control: Fresh buffalo dung with 2 cm depth. Different letters in each family indicate significantly different based on Mann-Whitney test p = 0.05.




Figure 4. Coverage of pioneer local Leguminosae/broadleaf plant, Cyperaceae, and Poaceae until 60 days after sowing on mulches

Note: A: Leguminosae/Broadleaf plant; B: Cyperaceae; C: Poaceae; B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; M₃: 0.2 cm depth; Control: Fresh buffalo dung with 2 cm depth.




Figure 5. Coverage of some pioneer local plant families

Note: A: Control plot; B: Hydroseeding treatment plot. Scale represents 2 cm.

Among the five species of Leguminosae family, namely *C. cajan*, *C. pallida*, *D. triflorum*, *I. spicata*, and *S. grandiflora*, there were only three species grown and had root nodules until 60 days after sowing. They were *C. cajan*, *C. pallida*, and *S. grandiflora* (Figure 6). It was different from a green house experiment conducted by Baiti and Arisoelaningsih (2015), which showed that growth of nodules on *C. pallida*, *D. triflorum*, and *I. spicata* after 37 days after sowing. The inside nodule was still yellow and not red, so that the nodule had not been functional. Number of functional root nodules could increase of soil nitrogen. Soil nitrogen fixation by Leguminosae increases the limited nitrogen level in post-mining soil for root growth (Singh et al., 2012). Post-mining soils generally have a low nitrogen content

that limits vegetation establishment and sustainable productivity (Tripathi and Singh, 2008).

Legume cultivation on marginal soils would be a multiple functions, such as cultivation crop, forage provision, green manure, and reduced artificial fertilizers use (Bruning and Rozema, 2013). The average number of three species nodules varied in each treatment. Root nodules number of *C. cajan*, *C. pallida*, and *S. grandiflora* in mulch with 4 cm depth (M_1) tended to be more than mulch with 6 cm depth (M_2) (Figure 6). Among three pioneer local plant families harvested on 60 days after sowing, Cyperaceae showed the highest value of root/shoot ratio compared to Leguminosae/broadleaf plant and Poaceae grown on all treatments (Figure 7).




Figure 6. Number of pioneer local Leguminosae root nodules until 60 days after sowing on mulches
 Note: Sg: *S. grandiflora*; Cc: *C. cajan*; Cp: *C. pallida*; B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; Control: Fresh buffalo dung with 2 cm depth. Different letters in each species indicate significantly different based on Duncan test p=0.05.




Figure 7. Root/shoot ratio value of three pioneer local plant families

Note: L: Leguminosae; C: Cyperaceae; P: Poaceae; B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; Control: Fresh buffalo dung with 2 cm depth. Different letters in each family indicate significantly different based on Duncan test p=0.05.

Although root/shoot ratio of Cyperaceae was higher than other two families, but Leguminosae root system was longer and developed better in the post-mining soil comparing to the shoot growth. Leguminosae had root/shoot ratio value <1 in B₁M₁, B₂M₁, B₂M₂, and control. Poaceae root/shoot ratio value >1 was found in B₁M₁, B₁M₂, B₂M₁, and B₂M₂ (Figure 7). This accorded to experiment conducted by Baiti (2015), that Leguminosae had root/shoot ratio <1 , and Rahma report (2015), which indicated that the root/shoot ratio >1 was found in Poaceae. Root/shoot ratio value >1 indicates that the plant root size is longer than the shoot. Some Poaceae showed a fast growth capacity, shallow rooting, and naturally able to creep well (Fan et al., 2013). Root/shoot ratio varied among species, growth rate, soil character, and available nutrients in the media. According to Tripathi et al. (2012), well developed root system had an important role in the rocks slope stabilization in coal mining areas. Root provided mechanical reinforcement to waste material through root proliferation. Species characteristic difference in growth and its

interactions also determine succession rate in unfavorable environmental conditions (Walker and del Moral 2008).

Productivity of some pioneer local plant families on mulches

The highest total plant productivity was found in control plot and reached 97 g/m²/month (Figure 8). Shoot productivity of all treatments was similar and greater value than the root one. Plant root and total productivity in hydroseeding plots were significantly lower than those of control plots. It revealed that the different mulches effected more significantly on the root part than on the shoot one.

Plant productivity was influenced by environmental factors, available nutrients in media, and density of growing seeds (Susilowati, 1999). Structural vegetation change in plant coverage and biomass production correlated with some soil changes such as accumulation of organic matter, nutrient content, porosity, and total nitrogen (Alday et al., 2012).




Figure 8. Plants productivity on mulches

Note: B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; Control: Fresh buffalo dung with 2 cm depth. Different letters indicate significantly different based on Mann-Whitney test $p=0.05$.


Multivariate profile of hydroseeding success

Based on multivariate analysis using biplot and clustering, it was shown that in the 1st application the artificial and natural hydroseeding had a


respective success and advantage (Figure 9A). There were three profiles based on germination characteristics, growth, and plants families (TL), high Poaceae germination (PG), total productivity (TP), coverage (Cov), and maximum height

(MH). The second profile, was B₁M₁, B₂M₁, and B₂M₂, that characterized by high Leguminosae (LG) and Cyperaceae germination (CG). While the B₁M₂ treatment was the third productivity. The first profile was control as the best mulch and showed by a long Leguminosae germination time (LT), leaves number of three profile and was characterized by a long Cyperaceae (CT) and Poaceae germination time (PT). Comparing to the

1st application, M₃ treatment showed a higher profile of shoot productivity (SP) and maximum height (MH) rather than the previous results treatments or control. In addition, its coverage (Cov) was similar to the profile of control and even greater than other treatments. It showed an important progress of hydroseeding medium for seedling growth (Figure 9B).



A



B

Figure 9. Interaction among seedling density, growth, and plant productivity of treatment mulches and control based on biplot analysis.

Note: A: 1st application; B: 2nd application; B₁: 360 seeds density; B₂: 720 seeds density; M₁: 4 cm depth; M₂: 6 cm depth; M₃: 0.2 cm depth; Control: Fresh buffalo dung with 2 cm depth; LG: Leguminosae germination; LT: Leguminosae germination time; CG: Cyperaceae germination; CT: Cyperaceae germination time; PG: Poaceae germination; PT: Poaceae germination time; Cov: coverage; TL: Leaves number of three plant families; MH: Maximum height; TP: Total productivity; SP: Shoot productivity.

In accordance with a study conducted by Samir et al. (2016), organic manure content in growing medium had significant effect on seed germination and seedling growth characteristics. A quite similar result reported by Ngo et al. (2011) indicated that the addition of buffalo manure content improved the organic matter quality of disturbed tropical soils, and this improvement positively affected plant growth. A comparison among soil, mixed media I (100% cow dung), and mixed media II (70% cow dung + 30% food industry sludge), performed by Yadav and Garg (2015), showed higher value of several physico-chemical characteristics, such as pH, electric conductivity, total organic carbon, total Kjeldhal nitrogen, total potassium, total phosphorus, and carbon nitrogen ratio, in both of mixed media rather than the soil. Therefore, the artificial hydroseeding using local seed M₃ was more optimal but still need to be improved by increasing seed viability and germination rate, by selecting seeds from more diverse fast growing species of three families, as well as by controlling mulch thickness. For better seed germination and plant growth, we focus to the seed quality and appropriate mulch composition. Besides, revegetation success should be monitored more than 3 months for ensuring higher plant coverage and biomass accumulation.

Conclusion

Based on the findings, it was concluded that the hydroseeding mulch could be served as a medium for pioneer local plant seeds germination and growth, especially for Leguminosae and comparable to buffalo dung. Percentage of Leguminosae seeds germination was higher than Cyperaceae and Poaceae seeds. Meanwhile, the growth of each family was specific depended on its respective habitus. Although root/shoot ratio of Cyperaceae was higher than other two families, but Leguminosae root was longer following to shoot growth. Hydroseeding plant productivity of 1st application was lower than control plots. In contrast, plant productivity of 2nd application was higher. Mulch with 720 seeds density (B₂) and 4 cm depth (M₁) was more optimal used in hydroseeding applications based on seed germination percentage of three pioneer local plant families, while mulch with 0.2 cm depth (M₃) was more effective for growth and productivity of plant.

Acknowledgement

The authors praised Allah SWT, so that the research and preparation of scientific work could be finished.

The authors also wished to thank the management of PT Amanah Anugerah Adi Mulia site Kintap and the entire staff of the SHE Department, who had provided land and facilities for supporting this research. Infinite gratitude was also extended to all JBUB lecturers, TU JBUB employees, Coal Mining Teams, and reviewers for all helps in completion of this research.

References

- Abella, S.C., Gunn, J.L., Daniels, M.L., Springer, J.D. and Nyoka, S.E. 2009. Using a diverse seed mix to establish native plants on a Sonoran Desert burn. *Native Plants Journal* 10 (1):21-31.
- Alday, J.G., Marrs, R.H. and Martinez-Ruiz, C. 2012. Soil and vegetation development during early succession on restored coal wastes: a six-year permanent plot study. *Plant and Soil* 353:305-320.
- Anshari, M.F. 2015. Revegetation of postmining soil using hydroseeding of local plant seeds in Tanah Laut Regency, South Kalimantan Province. Biology Department, Faculty of Mathematics and Natural Sciences, Brawijaya University, Malang. Thesis (*in Indonesian*).
- Baiti, R.N. 2015. Adaptation of some local Papilionaceae grown in coal postmining soil from South Kalimantan. Biology Department, Faculty of Mathematics and Natural Sciences, Brawijaya University, Malang. Thesis (*in Indonesian*).
- Baiti, R.N. and Arisoelaningsih, E. 2015. Reclamation of coal postmining soil using hydroseeding of local species of Papilionaceae beans. *Jurnal Biotropika* 3(1):36-41. (*in Indonesian*).
- Bruning, B. and Rozema, J. 2013. Symbiotic nitrogen fixation in legumes: perspectives for saline agriculture. *Environmental and Experimental Botany* 92:134-143.
- Darmawan, A. and Irawan, M.A. 2009. Reclamation of coal post-mining land in PT Berau Coal, East Kalimantan. Proceedings of science and technology workshop on forest rescue through coal postmining land rehabilitation. Research Center for Dipterocarps, Samarinda (*in Indonesian*).
- Decree of the Minister of Energy and Mineral Resources No. 1211.K/008/M.PE/1995 on prevention and mitigation of environmental destruction and pollution.
- Decree of the Minister of Forestry and Plantation No. 146/Kpts-II/1999 on guidelines for postmining reclamation in forest area.
- Djajakirana, G. 2001. Soil degradation affected by agricultural practices. Department of Soil Science and Land Resources, Faculty of Agricultural, Bogor Agricultural University, Bogor (*in Indonesian*).
- Fan, J., Huang, C., Yang, C., Liao, K. and Liao, W. 2013. Effect evaluation of shotcrete vegetation mulching technique applied to steep concrete-face slopes on a highway of Taiwan. *Paddy and Water Environment* 11:145-159.
- Faucete, L.B., Risse, L.M., Jordan, C.F., Cabrera, M.L., Coleman, D.C. and West, L.T. 2006. Vegetation and soil quality effects from hydroseed and compost blankets used for erosion control in

- construction activities. *Journal of Soil and Water Conservation* 61(6):355-361.
- General Directorate of Fiscal Balance at the Ministry of Finance of the Republic of Indonesia. 2012. Economic and finance study of South Kalimantan Province (in Indonesian).
- Hammer, Ø., Harper, D.A.T. and Ryan, P.D. 2001. PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4(1):1-9.
- Harun, N., Chaudhry, A.S., Shaheen, S., Ullah, K. and Khan, F. 2017. Ethnobotanical studies of fodder grass resources for ruminant animals, based on the traditional knowledge of indigenous communities in Central Punjab Pakistan. *Journal of Ethnobiology and Ethnomedicine* 13(1):56.
- Kak, A., Devi, L.C., Gupta, V. and Singh, N. 2007. Response of seed treatments on seed germination in wild *Crotalaria* species. *Acta Horticulturae* 752:261-265.
- Lancelotti, C. and Madella, M. 2012. The 'invisible' product: developing markers for identifying dung in archaeological contexts. *Journal of Archaeological Science* 39:953-963.
- Manaf, M.H. 2009. The environmental impacts of small-scale mining in Indonesia. www.gemeed.cl. Accessed on 4th December 2014 (in Indonesian).
- Ministry of Energy and Mineral Resources (ESDM). 2008. Center for data and information on energy and mineral resources. <http://www.esdm.go.id/> Accessed on 5th November 2014 (in Indonesian).
- Naikawadi, V.B. 2016. Seed germination of important medicinal plant *Tephrosia purpurea* (Linn.), Pers. *International Research Journal of Natural and Applied Sciences* 3(8):186-207.
- Ngo, P.T., Rumpel, C., Dignac, M., Billou, D., Duc, T.T. and Jouquet, P. 2011. Transformation of buffalo manure by composting or vermicomposting to rehabilitate degraded tropical soils. *Ecological Engineering* 37(2):269-276.
- Oliveira, G., Clemente, A., Nunes, A. and Correia, O. 2013. Limitations to recruitment of native species in hydroseeding mixtures. *Ecological Engineering* 57:18-26.
- Rahma, A.F. 2015. Productivity of local grasses grown by hydroseeding in monoculture and polyculture on post-coal mining soil from South Kalimantan. Biology Department, Faculty of Mathematics and Natural Sciences, Brawijaya University, Malang. Thesis (in Indonesian).
- Rochani, S. and Damayanti, R. 1997. Acid main drainage: general overview and strategies to control impacts. *Indonesian Mining Journal* 3(2):36-42.
- Samir, M., Sai, R. and Prasad, B. 2016. Effect of organic manures on seed germination and seedling growth of Khirni. *Indian Forester* 142(7):666-669.
- Singh, R.S., Tripathi, N. and Chauhya, S.K. 2012. Ecology study of revegetated coal mine spoil of an Indian dry tropical ecosystem along an age gradient. *Biodegradation* 23:837-849.
- Subandrio, A., Sukarman and Tambunan, R.T. 2009. Reclamation process in PT Adaro Indonesia. Proceedings of science and technology workshop on forest rescue through coal postmining land rehabilitation. Research Center for Dipterocarps, Samarinda (in Indonesian).
- Susilowati, A. 1999. Distribution and growth of *Scirpus litoralis* Schard. in mangrove forest of Resort Pemangku Hutan Cemara Indramayu. *Biosmart* 1:41-48 (in Indonesian).
- Tripathi, N. and Singh, R.S. 2008. Ecological restoration of mined-out areas of dry tropical environment, India. *Environmental Monitoring and Assessment* 146:325-337.
- Tripathi, N., Singh, R.S. and Chauhya, S.K. 2012. Dump stability and soil fertility of a coal mine spoil in Indian dry tropical environment: a long-term study. *Environmental Management* 50:695-706.
- Utomo, B. 2006. Seeds ecology. <http://repository.usu.ac.id>. Accessed on 29th June 2015 (in Indonesian).
- Walker, L.R. and del Moral, R. 2008. Lessons from primary succession for restoration of severely damaged habitats. *Applied Vegetation Science* 12:55-67.
- Walker, L.R. and del Moral, R. 2008. Transition dynamics in succession: implications for rates, trajectories, and restoration. In: Suding, K. and Hobbs, R.J. 2008. New models for ecosystem dynamics and restoration. Chapter 3 (eds). Island Press, Washington DC.
- Yadav, A. and Garg, V.K. 2015. Influence of vermicomposting on chickpea (*Cicer arietinum* L.) growth and photosynthesis pigments. *International Journal of Recycling of Organic Waste in Agriculture* 4:299-305.
- Zereen, A., Bokhari, T.Z. and Khan, Z. 2013. Ethnobotanical usages of grasses in Central Punjab-Pakistan. *International Journal of Scientific and Engineering Research* 4(9):452-461.
- Zipper, C.E., Burger, J.A., Barton, C.D. and Skousen, J.G. 2012. Rebuilding soils on mined land for native forests in Appalachia. *Soil Science Society of America Journal* 77:337-349.